

**Cytotoxic activity of thirteen endemic and rare plants from Chaharmahal and Bakhtiari
Province in Iran**

Seyed Ahmad Emami^a, Shima Khashami^c, Elham Ramazani^b, Maryam Akaberi^a, Milad
Iranshahi^a, Seyed Mahammad Kazemi^c and Zahra Tayarani-Najaran^{d*}

^a*Department of Pharmacognosy, School of Pharmacy, Mashhad University of Medical Sciences,
Mashhad, Iran.* ^b*Department of Biology, Faculty of Science, Ferdowsi University of Mashhad,
Mashhad, Iran.* ^c*Medical Toxicology Research Center, Mashhad University of Medical Sciences,
Mashhad, Iran.* ^d*Biotechnology Research Center, Mashhad University of Medical Sciences,
Mashhad, Iran.*

E-mail address: tayaraninz@mums.ac.ir

Abstract

Chaharmahal and Bakhtiari Province is a most important endemism states of flora of Iran with a considerable plant species diversity in the country. In the present study the cytotoxic activity of 13 plant species grown in Chaharmahal and Bakhtiari have been

evaluated on prostate (PC-3), breast (MCF-7), liver (HepG2), ovary (CHO) and melanoma (B16-F10) cancer cell lines. The cytotoxicity and apoptotic activity of methanol extracts were evaluated using resazurin reagent and flow cytometry of PI stained cells, respectively. Methanol extracts of *Dionysia sawyeri*, *Stachys obtusirena* and *Cicer oxyodon* on CHO cell line ($p < 0.05$) and *D. sawyer* and *L. album* on B16/F10 cell line ($p < 0.05$) showed significant cytotoxic effects and increased apoptosis. It is generally suggested that plant species with low IC₅₀ values are likely to be used as anti-cancer compounds in reducing cancer progression in scientific studies.

Keywords: *Achillea*, *Ajuga*, *Aristolochia*, *Cicer*, *Dianthus*, *Dionysia*, *Linum*, *Nepeta*, *Phlomis*, *Picris*, *Scutellaria*, *Stachys*, *Tanacetum*, Cytotoxic, Apoptosis.

Introduction

Cancer is the most recognized term used for more than 100 different types of malignancies that can affect the body (1). The number of various cancer deaths in 2015 was 8.8 million people which is equivalent 1 in 6 worldwide death cases (2). One of the common problems in treating cancers is the tumor cells resistance to conventional chemotherapeutic drugs. This phenomenon leads to formation of cells with more aggressive phenotype which are more likely to metastases to other tissues (3). Additional to the mentioned problems in treating cancer, side effects of chemotherapy also attract researchers to investigate for new approaches in cancer treatments such as the use of natural resources as therapeutic compounds (4).

In the present study we have chosen 13 different endemic and rare plant species from Chaharmahal and Bakhtiari Province, Iran, which there is not any report about the cytotoxic activity of plant. Plants from the same genus have some similarities in case of phytochemicals and consequently in their biologic activity, therefore the results of studies on same species discussed. Lu and colleagues (2016) reported that, the phlomisioside from the plant *Phlomis younghusbandii* have a significant inhibitory effect on the growth, proliferation, migration and invasive properties of A549 (human lung cancer cells) cancer cell line with IC₅₀ of 54.51 μ M and induces apoptosis (5). Ji and his colleagues (2015) evaluated cytotoxic activity of *Scutellaria baicalensis* on HepG2 (human liver cancer cells), SW480 (Human colon cancer cells) and MCF-7 (human breast adenocarcinoma cell) cancer cell lines. Results showed that most of the flavones exhibited a significant cytotoxic effect (6). Tayarani-Najaran (2010) and colleagues addressed the cytotoxic effects of total methanol extract and different fractions of *Scutellaria lindbergii* on AGS (Human stomach cancer cells), HeLa (Human cervix cancer cells), MCF-7 and PC12 (Rat adrenal gland cancer cells) cells. Based on the results, methylene chloride fraction has shown the most potent cytotoxic activity among other fractions and decreased cell viability (7). *Scutellaria pinnatifida* and its active component neobaicalein (skullcapflavone II) and wogonin showed strong cytotoxic activity against HL-60 and K562 leukemic cell lines (8).

Due to specific climatic conditions and the high diversity of plant species in different regions of Iran, investigation on the therapeutic properties and molecular functional mechanisms of the rare plant species for exploring new drugs for treating various diseases, including cancer is worthy. One of the rich areas with diverse plant

species in Iran is Chaharmahal and Bakhtiari Province. The Chaharmahal and Bakhtiari Province is located in the middle of the mountains of the west of Iran and the plain of Isfahan. The situated area is located between 31 14` and 33 47` N (latitude), 49 49` and 51 34` E (longitude). About 1400000 hectares of the total area of the Province which is equivalent to 86.6% of all total area is occupied with forests and pastures (9).

In this study cytotoxic effects and apoptosis induction of methanol extract from thirteen rare plants from Chaharmahal and Bakhtiari Province, including species from the families listed in Table 1, were assessed on the human prostate cancer (PC-3), (MCF-7), (HepG2), Chinese hamster ovary cells (CHO) and murine melanoma (B16-F10) cancer cell lines. To the best our knowledge this is the first report on cytotoxic activity of the plants.

Table 1. Medicinal plants evaluated for cytotoxic activity from Chaharmahal and Bakhtiari Province of Iran.

Species	State	Family	Voucher number	Location
<i>Achillea kellalensis</i> Boiss. & Hausskn.	endemic	Asteraceae	13206	Gelougerd, the Northern slopes of mountain Kalar
<i>Ajuga chamaecistus</i> Ging. ex Benth.	rare	Lamiaceae	13201	Shahrekord, the mountain Farhangian
<i>Aristolochia olivieri</i> Colleg. ex Boiss.	endemic	Aristolochiaceae	13202	Malkhalifeh, Shirmard village
<i>Cicer oxyodon</i> Boiss. & Hohen.	rare	Fabaceae	13207	Malkhalifeh, Shirmard village
<i>Dianthus orientalis</i> Adams	rare	Caryophyllaceae	13208	Hafshejan, Jouneghan
<i>Dionysia sawyeri</i> (Watt) Wendelbo	endemic	Primulaceae	13205	Malkhalifeh, Shirmard village
<i>Linum album</i>	endemic	Linaceae	13204	Shahrekord, the mountain

Kotschy ex Boiss.				Farhangian
<i>Nepeta glomerulosa</i> Boiss.	endemic	Lamiaceae	13200	Shahrekord, castle Gharak
<i>Phlomis aucheri</i> Boiss.	endemic	Lamiaceae	13199	Shahrekord, castle Gharak
<i>Picris strigosa</i> M. Bieb.	rare	Asteraceae	13203	Malkhalifeh, Shirmard village
<i>Scutellaria multicaulis</i> Boiss.	endemic	Lamiaceae	13198	Malkhalifeh, Shirmard village
<i>Stachys obtusicrena</i> Boiss.	endemic	Lamiaceae	13196	Avargan, the mountain Kalar
<i>Tanacetum dumosum</i> Boiss.	endemic	Asteraceae	13197	Malkhalifeh, Shirmard village

Experimental

Plant materials

Thirteen species of endemic and rare plants from different families were collected in spring and summer 2105 from various regions of Chaharmahal and Bakhtiari Province, southwestern of Iran and identified by H. A. Shirmardi (Table 1). Voucher specimens of the species were deposited in the herbarium of School of Pharmacy, Mashhad University of Medical Sciences, Iran.

Preparation of extracts

The aerial parts of each species dried in shadow and powdered. Then, 100 g of each powder was macerated with methanol for 24 hr. at controlled room temperature (25°C). The macerated powder sample was percolated using pure methanol. Then the methanol extracts were concentrated via a rotary evaporator under reduced pressure at 50°C and subsequently freeze dried. All extracts were stored at -20 °C. The yield percentage of the obtained extracts were presented in Table 2.

Table 2. The extraction yield % of medicinal plants.

Species	Extraction yield %
<i>Achillea kellalensis</i>	11.77%
<i>Ajuga chamaecistus</i>	16.44%
<i>Aristolochia olivieri</i>	19.25%
<i>Cicer oxyodon</i>	16.46%
<i>Diantus orientalis</i>	8.31%
<i>Dionysia sawyeri</i>	1.84%
<i>Linum album</i>	14.6%
<i>Nepate glomerulosa</i>	7.6%
<i>Phlomis aucheria</i>	16%
<i>Picris strigosa</i>	4.4%
<i>Scutellaria multicaulis</i>	7.41%
<i>Stachys obtusicrena</i>	17.01%
<i>Tanaetum dumosum</i>	11.37%

Table 3. Biological properties and chemical constituents of medicinal plants from Chaharmahal and Bakhtiari Province of Iran.

Species	Chemical constituents	Biological activities
<i>Achillea kellalensis</i>	Camphor (34.0%), borneol (12.6%), β -thujone (12.5%), 1,8-cineole (11.3%), bornyl acetate (7.3%), camphene (7.0%) (Rustaiyan et al., 1999)	Antioxidant (Pirbalouti et al., 2010a), antibacterial (Gharibi et al., 2013)

<i>Ajuga chamaecistus</i>	Melilotoside, phenylethyl glycosides, phytoecdysteroids (Sadati et al., 2012)	Antidiabetic (Khanavi et al., 2014), anti-inflammatory (Eskandani et al., 2016), antibacterial (Moshefi et al., 2014)
<i>Aristolochia olivieri</i>	-	-
<i>Cicer oxyodon</i>	-	-
<i>Dianthus orientalis</i>	-	-
<i>Dionysia sawyeri</i>	-	-
<i>Linum album</i>	Podophyllotoxin, 5-methoxypodophyllotoxin (Smollny et al., 1998)	Antitumor (Weiss et al., 1975)
<i>Nepeta glomerulosa</i>	Geranyl acetate (17.0%), limonene (12.0%), eucalypto (5.8%), bornyl acetate (5.3%), citronellal (4.9%), spathulanol (4.2%), sabinene (3.9%), β -ocimene (3.9%), β -sesquiphellandrene (2.8%), neryl acetate (2.5%), α -humulene (2.4%), α -pinene (2.3%), humulene oxide (2.2%), norsolanadione (2.1%), terpinen-4-ol (2.0%) (Nezhadali et al., 2011)	Antibacterial (Nezhadali et al., 2011)
<i>Phlomis aucherii</i>	E-anethole (24.58%), germacrene D (11.1%), bicyclogermacrene (6.3%), spathulenol (6.01%), bicyclogermacrene (5.58%), nerylacetate (4.58%), and germacrene B (4.53%) (Taherkhani, 2015)	Antioxidant, antibacterial (Taherkhani, 2015)
<i>Picris strigosa</i>	-	-
<i>Scutellaria multicaulis</i>	Trans-caryophyllene (34.6%), caryophyllene oxide (12.2%), linalool (10.7%), germacrene D (5.5%) (Asadollahzadeh and Rajaie, 2014)	Antioxidant (Souria et al., 2004)
<i>Stachys obtusicrena</i>	α -pinene (34.6%), germacrene D (8.0%), bicyclogermacrene (7.8%) (Jamzad et al., 2009)	Antibacterial (Masoudi et al., 2012), Anti-inflammatory (Amirghofran, 2010), Antimicrobial (Javidnia et al., 2009)
<i>Tanacetum dumosum</i>	Borneol (27.9%), bornyl acetate (18.4%), 1,8-cineol (17.5%), α -terpineol (5.3%), cis-chrysanthenyl acetate (3.3%), camphene (2.7%), terpinene-4-ol (1.9%) (Ghanbarian et al., 2015)	-

Cell culture and Treatment

The prostate (PC-3), breast (MCF-7), liver (Hep G2), ovary (CHO) and melanoma (B16/F10) cancer cell lines (code numbers: C427, C135, C158, C111 and C540) were obtained from Cell Bank at the Pasteur Institute (Tehran, Iran). CHO were cultured in F-12K medium (Sigma) with 10% (v/v) fetal bovine serum, 100U/mL penicillin and 100 mg/mL streptomycin. B16 F10 and MCF7 were cultured Dulbecco's Modified Eagle's Medium (DMEM) medium (Sigma) with 10% (v/v) fetal bovine serum, 100U/mL penicillin and 100 mg/mL streptomycin. While other cell lines were cultured in RPMI 1640 medium (Sigma) with 10% (v/v) fetal bovine serum, 100U/mL penicillin and 100 mg/mL streptomycin. Then all cells were kept at 37 °C in a humidified atmosphere (90%) containing 5% CO₂. For each concentration and time course study, there was a control sample, which remained untreated and received an equal volume of the solvent (10).

Cell Viability

The resazurin reagent is a cell viability indicator that allow to measure cytotoxicity of various chemicals components. For detecting cell viability, all of cancer cell lines (104 cells per well) were seeded in 96-well plates and were incubated with the methanol extract of each species (50 and 100 μ M) for 48 h. Then resazurin reagent (20 μ M) was added to each well and incubated for 4 h. The cell viability was assessed at absorbance of 600 nm with ELISA micro plate reader (Awareness, Palm City, FL, USA) (11).

Table 4. Cytotoxicity (% of viability) of methanol extract of medicinal plants from Chaharmahal and Bakhtiari Province of Iran.

Cell line→	Concentration ¹														
	PC3			MCF7			HepG2			CHO			B16-F10		
Plant name↓	0	50	100	0	50	100	0	50	100	0	50	100	0	50	100
<i>Achillea kellalensis</i>	100.0±1.9.3	107.0±2.9.0	104.3±3.2	100.0±3.9.7	99.7±27.6	110.2±4.5.5	99.9±13.5	87.5±12.9	93.6±10.5	100.0±3.9.3	72.1±23.5	77.7±19.5	100.0±7.0	110.6±8.4	88.7±19.1
<i>Ajuga chamaecistus</i>	100.0±1.9.3	105.4±3.3.3	100.8±4.1.5	100.0±3.9.7	105.9±31.5	96.8±38.8	99.9±13.5	103.5±9.3	98.0±11.3	100.0±3.9.3	66.7±15.7	54.0±8.9	100.0±7.0	102.0±15.2	99.9±8.9
<i>Aristolochia olivieri</i>	100.0±1.9.3	105.7±2.1.0	111.0±1.9.7	100.0±3.9.7	111.3±9.8	121.4±5.6	99.9±13.5	108.1±9.2	112.0±6.1	100.0±3.9.3	90.1±20.6	79.2±30.6	100.0±7.0	98.5±13.8	108.8±45.1
<i>Cicer oxyodon</i>	100.0±1.9.3	107.7±3.9.5	101.5±2.2.9	100.0±3.9.7	101.0±36.3	96.8±37.5	99.9±13.5	113.0±6.6	120.6±8.4	100.0±3.9.3	63.3±20.7*	68.5±28.2*	100.0±7.0	81.0±7.8	71.7±10.1
<i>Dianthus orientalis</i>	100.0±1.9.3	107.6±2.2.7	105.8±2.3.6	100.0±3.9.7	104.9±26.1	97.5±39.2	99.9±13.5	89.4±9.5	95.4±10.9	100.0±3.9.3	73.4±23.0	79.7±24.6	100.0±7.0	81.0±30.6	102.3±18.4
<i>Dionysia sawyeri</i>	100.0±1.9.3	95.3±27.8	98.6±29.0	100.0±3.9.7	91.2±13.7	104.6±1.4.4	99.9±13.5	93.0±7.2	87.3±6.0	100.0±3.9.3	62.3±16.8*	58.6±19.7*	100.0±7.0	76.5±21.3*	62.9±12.8*
Doxorubicin	100.0±1.9.3	58.6±18.7*	46.6±17.1*	100.0±3.9.7	36.4±26.8**	52.7±21.1*	99.9±13.5	38.8±15.2**	40.0±15.2*	100.0±3.9.3	64.8±26.3*	71.2±15.7*	100.0±7.0	20.4±12.5**	36.2±11.7**
<i>Linum album</i>	100.0±1.9.3	110.8±2.0.7	100.9±1.5.1	100.0±3.9.7	120.3±21.4	120.0±1.3.3	99.9±13.5	96.2±10.2	97.7±13.6	100.0±3.9.3	88.1±16.8	74.7±11.7	100.0±7.0	63.6±6.8*	69.6±20.1*
<i>Nepeta glomerulosa</i>	100.0±1.9.3	102.4±2.0.4	106.1±1.9.5	100.0±3.9.7	125.9±13.7	117.8±1.2.9	99.9±13.5	96.7±10.3	101.5±8.0	100.0±3.9.3	88.9±29.4	86.5±28.3	100.0±7.0	102.5±10.4	111.8±13.8
<i>Phlomis aucheri</i>	100.0±1.9.3	103.8±2.7.0	97.8±12.8	100.0±3.9.7	112.7±12.5	116.7±1.6.6	99.9±13.5	89.7±10.0	97.6±11.2	100.0±3.9.3	98.2±19.9	75.8±24.6	100.0±7.0	104.0±16.9	96.3±21.0
<i>Picris strigosa</i>	100.0±1.9.3	106.4±2.3.0	104.5±3.6.0	100.0±3.9.7	114.0±11.1	112.0±1.1.1	99.9±13.5	102.5±8.9	105.1±5.9	100.0±3.9.3	70.8±17.8	76.0±18.4	100.0±7.0	114.5±9.5	92.9±21.9

<i>Scutellaria multicaulis</i>	100.0±1.9.3	96.8±25.2	96.7±36.1	100.0±3.9.7	104.8±26.1	91.9±5.8	99.9±13.5	92.5±8.6	94.8±4.2	100.0±3.9.3	58.5±43.9	72.6±42.0	100.0±7.0	89.3±16.5	79.5±20.8
<i>Stachys obtusirena</i>	100.0±1.9.3	120.2±1.8.2	98.3±21.0	100.0±3.9.7	102.8±25.7	109.8±1.0.4	99.9±13.5	90.2±12.8	87.5±5.3	100.0±3.9.3	61.1±20.0*	53.7±6.3*	100.0±7.0	86.7±21.3	98.7±17.7
<i>Tanacetum dumosum</i>	100.0±1.9.3	91.6±29.8	92.2±28.2	100.0±3.9.7	79.3±35.4	100.0±1.4.8	99.9±13.5	99.7±7.8	112.3±8.3	100.0±3.9.3	89.9±29.8	78.8±32.3	100.0±7.0	87.0±25.9	74.2±38.2

¹ The concentration of extracts is presented in µg/mL.

Dunnet's Multiple Comparison Test; * $p < 0.05$, ** $p < 0.001$

Flow cytometry analysis of Apoptosis

Flow cytometry and PI staining of treated cells to detect a sub-G1 peak evaluated Apoptotic cells. CHO cells (105 cells per well) were cultured into 24-well plates and were treated with the methanol extract of *D. sawyeri*, *S. obtusirena* and *C. oxyodon* (50 and 100 µM) for 48 h and also B16/F10 were treated with the methanol extract of for 48 h. Cells were washed with phosphate-buffer saline (PBS). After trypsinization cells were harvested and incubated at 4°C in dark with 400 µL of hypotonic buffer (50 µg/mL PI in 0.1% sodium citrate and 0.1% Triton X-100) for 30 s before flow cytometric analysis using a Partec flow cytometer (GmbH, Münster, Germany) (11).




Figure 1. Flow cytometry histograms of apoptosis assays by PI method of CHO and B16/F10 cells: (A) CHO cells were incubated with 50, 100 μ M of Methanol extracts of *Dionysia sawyer*, *Stachys obtusirena* and *Cicer oxyodon* and (B) B16/F10 cells were incubated with 50, 100 μ M of *D. sawyer* and *Linnum album* for 48 h. All components induced cell death through apoptosis. All experiments were done in triplicate.

Statistics Analysis

One-way analysis of variance (ANOVA) and Turkeys-Kramer post hoc were used for data analysis. All results were expressed as mean \pm SD and p-values below <0.05 was regarded statistically significant.

Results

Based on the results, the extracts of *D. sawyeri*, *S. obtusirena* and *C. oxyodon* on CHO cell line ($p <0.05$) and the extracts of *D. sawyeri* and *L. album* on B16-F10 cell line ($p <0.05$) decreased cell viability and showed significant cytotoxic effects. (Tables 3 and 4). Also, results demonstrated that extracts of *D. sawyeri*, *S. obtusirena* and *C. oxyodon* on CHO cell line induced cell death through apoptosis (Fig. 1).

Discussion

Variation in plant species also the presence of exclusive plant species that have not been studied so far require extensive biological screening to find putative natural substances as effective agents for the treatment of disease. Many substances including natural products and phytochemicals have been screened to find an optimal treatment for cancer as the second cause of death worldwide.

In this study, the cytotoxic effects of methanol extract from 13 plant species from Chaharmahal and Bakhtiari Province investigated on PC-3, CHO, B16/F10, HepG2 and MCF-7 cancer cell lines. According to the results, methanol extract of *D. sawyeri*, *S. obtusicrena* and *C. oxyodon* in CHO cells and methanol extract of *D. sawyer* and *L. album* in the B16/F10 cells decreased cell viability and showed significant cytotoxic activity. Also methanol extract of *D. sawyeri*, *S. obtusicrena* and *C. oxyodon* increased apoptosis induction in CHO cells. This is the first study that evaluated the cytotoxic effects of rare plants from Chaharmahal and Bakhtiari Province. Regarding the novelty of the present study, there is not any similar evaluation on the same species have been investigated here. Since plants belonging to the same genus have similarities in presence of alike phytochemicals, we have searched for some evidence of cytotoxicity in similar species in the same genus.

Dionysia termeana is one of the similar species to *D. sawyeri*, inhibited the growth and proliferation of leukemia (K562) and lung carcinoma (A549) cell line with an IC50 less than 20 µg/ml by MTT staining and flow cytometry analyzes (12). *Dionysia termeana*, also significantly inhibited the growth and proliferation of lymphocyte cells (13). In our study, *D. sawyeri* exerted cytotoxic effect through decreasing cell viability and increasing amount of apoptosis.

There are many reports on the cytotoxic activity of the plants belongs to the genus *Stachys*. Jassbi et al. (2014) examined the cytotoxic effects of methanol and dichloromethane extracts of nine different species of *Stachys* on HL-60, K562 and MCF-7 cancer cells. The authors reported that dichloromethane extract of *S. pilifera* had the lowest IC50 on HL-60 (Human leukemia cancer cells), K562 (Human leukemia

cancer cells) and MCF-7 cancer cells (ranging from 33.1 to 18.4 µg/ml) (14). In another study it has been shown that *S. alopecuross* inhibit the growth of A375 (Human melanoma cancer cells), HCT116 (Human colon cancer cells) and MDA-MB 231 (Human breast cancer cells) cells with IC50 less than 20 µg/ml (15). Also, the volatile oil of *S. rupestris* has been shown to inhibit the growth and proliferation of PC-3 and MCF-7 cell lines (16). In our study, *S. obtusicrena* showed inducing effect on apoptosis and cell growth inhibition.

Cicer microphylluma similar species from the same genus of *C. oxyodon*, has been shown potent cytotoxic activity against mammary melanoma cell lines and human epidermis carcinoma in Dar et al., (2015) study which was attributed to the presence of luteolinin the plant (17). Isoflavones extracted from *C. arietinum* promoted the growth of MCF-7 cell line at low concentrations and inhibited the growth and proliferation of the cells at high concentrations (more than 1 mg/l) (18). Isoflavones extracted from the *C. arietinum* inhibited the growth and proliferation of two human breast cancer cell lines including SKBR3 (Human breast cancer cells) and MCF-7 (19).

Dionysia sawyeri, *S. obtusicrena*, *C. oxyodon* and *L. album* as the most cytotoxic and CHO and B16/F10 cells as the most sensitive cells were chosen for future mechanistic activity. In our study, *D. sawyeri*, *S. obtusicrena*, *C. oxyodon* increased apoptosis induction which was confirmed after PI staining of cells and flow cytometry analysis on CHO cells. In the present study for the first time, the cytotoxic activity of *L. album* was reported. *L. album* caused a dose-dependent cytotoxic activity on the B16-F10 cell line with minimal effect on other cells.

Conclusion

Two common types of cancer in young adults and young adult's female are melanoma and ovarian cancer. Melanoma is one of the most hazardous forms of skin cancer and which occurred in 232,000 people and resulted in 55,000 deaths in 2012. Surgery is mostly used to treat this cancer via removing involved parts. Ovarian cancer is the seventh most common cancer in women and the eighth leading cause of cancer death in the world. This occurred around 239,000 cases and resulted in 152,000 deaths worldwide in 2012. Treatment usually involves a combination of surgery, radiation therapy and chemotherapy.

Among four cytotoxic plant introduced in this study, Main constituents from *S. obtusirena* and *L. album* have been reported in table 3. Based on the diversity of the chemicals present in plants, it is suggested that further analytical and mechanistic evaluation supports the use of plant as potential anticancer agents.

Acknowledgement

Financial support of this study was provided by research affairs of Mashhad University of Medical Sciences by grant no: 931055 and performed as a part of a Pharm.D. thesis.

References

1. Stjernswärd J, Colleau SM, Ventafridda V. The World Health Organization cancer pain and palliative care program past, present, and future. *J. Pain Symptom Manage.* (1996) 12: 65-72.
2. <http://www.who.int>, 2016. 10 Facts on Cancer.
3. Simstein R, Burow M, Parker A, Weldon C, Beckman B. Apoptosis, chemoresistance, and breast cancer: insights from the MCF-7 cell model system. *Exp. Biol. Med.* (2003) 228: 995-1003.
4. Scheck AC, Perry K, Hank NC, Clark WD. Anticancer activity of extracts derived from the mature roots of *Scutellaria baicalensis* on human malignant brain tumor cells. *BMC Complement. Altern. Med.* (2006) 6: 27-36.
5. Lu XX, Ji XX, Bao J, Li QQ, Ji DD, Luo L. Inhibition of proliferation, migration and invasion of human non-small cell lung cancer cell line A549 by phlomisioside F from *Phlomis younghusbandii* Mukerjee. *Trop. J. Pharm. Res.* (2016) 15: 1413-1421.
6. Ji S, Li R, Wang Q, Miao WJ, Li ZW, Si LL, Qiao X, Yu SW, Zhou DM, Ye M. Anti-H1N1 virus, cytotoxic and Nrf2 activation activities of chemical constituents from *Scutellaria baicalensis*. *J. Ethnopharmacol.* (2015) 176: 475-484.
7. Tayarani-Najaran Z, Mousavi S, Asili J, Emami SA. Growth-inhibitory effect of *Scutellaria lindbergii* in human cancer cell lines. *Food Chem. Toxicol.* (2010) 48: 599-604.
8. Boozari M, Mohammadi A, Asili J, Emami SA, Tayarani-Najaran Z. Growthinhibition and apoptosis induction by *Scutellaria pinnatifida* A. Ham. on HL-60and K562 leukemic cell lines. *Environ. Toxicol. Pharmacol.* (2015) 39: 307-12.
9. Ezati Asar M, Varehzardi R, Rajabi Vasokolaei G, Haghi M, Fazelipor M. Regional disparities in the distribution of healthcare workers: evidence from Iran, Chaharmahal and Bakhtiari Province. *Glob. J. Health Sci.* (2015) 7: 374-378.

10. Mousavi SH, Motaez M, Zamiri-Akhlaghi A, Emami SA, Tayarani-Najaran Z. In-vitro evaluation of cytotoxic and apoptogenic properties of *Sophora pachycarpa*. Iran. J. Pharm. Res. (2014) 13: 665-673.
11. Tayarani-Najaran Z, Sareban M, Gholami A, Emami SA, Mojarrab M. Cytotoxic and apoptotic effects of different extracts of *Artemisia turanica* Krasch. on K562 and HL-60 cell lines. Sci. World J. (2013) 2013.
12. Amirghofran Z, Bahmani M, Azadmehr A, Ashouri E, Javidnia K. Antitumor activity and apoptosis induction in human cancer cell lines by *Dionysia termeana*. Cancer Invest. (2007) 25: 550-554.
13. Amirghofran Z, Bahmani M, Azadmehr A, Javidnia K, Miri R. Immunomodulatory activities of various medicinal herb extracts: effects on human lymphocytes apoptosis. Immunol. Invest. (2009) 38: 181-192.
14. Jassbi AR, Miri R, Asadollahi M, Javanmardi N, Firuzi O. Cytotoxic, antioxidant and antimicrobial effects of nine species of woundwort (*Stachys*) herbs. Pharm. Biol. (2014) 52: 62-67.
15. Venditti A, Bianco A, Nicoletti M, Quassinti L, Bramucci M, Lupidi G, Vitali LA, Petrelli D, Papa F, Vittori S. Phytochemical analysis, biological evaluation and micromorphological study of *Stachys alopecuroides* (L.) Benth. subsp. *divulsa* (Ten.) Grande endemic to central Apennines, Italy. Fitoterapia. (2013) 90: 94-103.
16. Erdogan EA, Everest A, De Martino L, Mancini E, Festa M, De Feo V. Chemical composition and *in-vitro* cytotoxic activity of the essential oils of *Stachys rupestris* and *Salvia heldreichiana*, two endemic herbs of Turkey. Nat. Prod. Commun. (2013) 8: 1637-1640.
17. Dar AA, Rath SK, Qaudri A, Singh B, Tasduq SA, Kumar A, Sangwan PL. Isolation, cytotoxic evaluation, and simultaneous quantification of eight bioactive secondary metabolites from *Cicer microphyllum* by high-performance thin-layer chromatography. J. Sep. Sci. (2015) 38: 4021-4028.
18. HaiRong M, HuaBo W, Zhen C, Yi Y, ZhengHua W, Madina H, Xu C, Haji Akber A. The estrogenic activity of isoflavones extracted from chickpea *Cicer arietinum* L. sprouts *in-vitro*. Phytother. Res. (2013) 27: 1237-1242.

19. Chen H, Ma, HR, Gao YH, Zhang X, Habasi M, Hu R, Aisa HA. Isoflavones extracted from chickpea *Cicer arietinum* L. sprouts induce mitochondria-dependent apoptosis in human breast cancer cells. *Phytother. Res.* (2015) 29: 210-219.