

Original Article

Investigation of Cytotoxic Activity in Four *Stachys* Species from Iran

Mahnaz Khanavi^a, Azadeh Manayi^a, Mahnaz Lotfi^b, Rofeyde Abbasi^b,
Maryam Majdzadeh^a and Seyed Nasser Ostad^{c*}

^aDepartment of Pharmacognosy, Faculty of Pharmacy and Traditional Iranian Medicines and Pharmacy Research Center, Tehran University of Medical Sciences, Tehran, Iran.

^bPharmaceutical Sciences Branch, Islamic Azad University, Tehran, Iran. ^cDepartment of Toxicology and Pharmacology, Faculty of Pharmacy and Pharmaceutical Sciences Research Center, Tehran University of Medical Sciences, Tehran, Iran.

Abstract

The aerial parts of *Stachys laxa* Boiss. and Buhse. from Siah-bishe in Mazandaran province, *Stachys trinervis* Aitch. and Hemsl. from Karaj in Alborz province, *Stachys subaphylla* Rech. F. and *Stachys turcomanica* Trautv. from Golestan province have been collected in May 2008. Total extracts were obtained through MeOH/H₂O (80/20) and then partitioned between CHCl₃, EtOAc and MeOH. These fractions and total extracts have been investigated for *in-vitro* cytotoxic activity against the colon carcinoma (HT-29), colorectal adenocarcinoma (Caco-2), breast ductal carcinoma (T47D) and Swiss mouse embryo fibroblast (NIH 3T3) cell lines using MTT assay (3-(4,5-di methyl thiazol-2-yl)-2,5-di phenyltetrazolium bromide). At each cell line, doses of 3.125, 6.25, 12.5, 25, 100, 200, 400 and 800 µg/mL in 1% (v/v) DMSO of all samples were tested. Ethyl acetate and chloroform fractions of *Stachys laxa* against proliferation of T47D and HT-29 cell lines and chloroform fraction of *Stachys subaphylla* and *Stachys subaphylla* ethyl acetate fraction toward T47D cell line exhibited highest cytotoxic activity (IC₅₀ < 50 µg/mL). Ethyl acetate and chloroform fractions of *Stachys turcomanica* against HT-29 cell line, except methanol fraction of *Stachys subaphylla*, the other extracts on T47D cell line, represented moderate cytotoxic activity (IC₅₀ < 70 µg/mL). All fractions of *S. trinervis* demonstrated no effective cytotoxic activity. IC₅₀ values confirmed that the growth and proliferation of HT-29 and T47D cells were most affected by chloroform and ethyl acetate fractions of *Stachys laxa* and *Stachys turcomanica* due to their nonpolar compounds.

Keywords: Cytotoxic activity; *Stachys laxa* Boiss. and Buhse.; *Stachys turcomanica* Trautv.; *Stachys subaphylla* Rech. F.; *Stachys trinervis* Aitch. and Hemsl.; MTT assay.

Introduction

The genus *Stachys* belongs to the plant family of Lamiaceae. The most species of this genus has been previously analyzed in numerous studies concerning their chemical composition,

pharmacological properties and therapeutic uses. This family is well represented in the flora of Iran, at least with 200-300 species in the world (1) and 34 species in Iran (2). Phytochemical investigation of some *Stachys* species has demonstrated phenolic acids, tannins (3, 4), flavonoids (5) and phenyl ethanoid glycosides (6, 7). There are some reports about pharmacological activities of this genus including anticancer (8,

* Corresponding author:

E-mail: ostadnas@sina.tums.ac.ir

9), antibacterial (10-13), antioxidant effects (14-16), anti-inflammatory (17-22), anti-nephritic (23) and anti-anxiety (24). Some *Stachys* species are used in folk medicine for healing wounds, disinfectant, treating abdominal pains, asthma, rheumatic and inflammatory disorders, anti-spasmodic and anti-fever (20, 25). Essential oils of *Stachys cretica* ssp. *lesbiaca* Rech. fil. and *S. cretica* ssp. *trapezuntica* Rech. fil. inhibit the growth of HL-60 and Ishikawa human tumor cell lines; the main component in both of them was germacrene D (26). There was an investigation about the essential oil of *S. turcomanica*; the major constituents were identified as germacrene D (17.4%), 7-epi- α -selinene (10.5%), β -elemene (9.2%) and β -pinene (8.6%) (27). Major components of *S. trinervis* oil were identified as α -pinene (42.68%), δ -2-carene (31.90%), 1,8-cineole (7.03%), limonene (4.39%) and (Z)- β -ocimene (4.21%). *S. subaphylla* oil major constituents were δ -2-carene (23.93%), α -pinene (19.29%), sabinene (19.11%), δ -3-carene (9.22%) and (Z)- β -ocimene (5.90%) (28). Thirty three constituents from essential oil of *S. laxa* have been identified and the major constituents were germacrene D (40.1%), β -caryophyllene (16.7%), β -phellandrene (5.5%), caryophyllene oxide (4.6%), linalool (3.2%) and α -cadinol (2.6%) (29). The cytotoxicity of some *Stachys* species against A431, HeLa and MCF-7 were examined. *S. recta* and *S. palustris* stem extract inhibit the growth of HeLa cells. An *S. rectum* was significantly active toward the breast MCF7 cell line (30). More than half of drugs in cancer therapy were obtained from natural products or related to them (31). Hence in this study, cytotoxic activity of *S. laxa* Boiss. and Buhse., *S. turcomanica* Trautv., *S. subaphylla* Rech. F., *S. trinervis* Aitch. and Hemsl. have been investigated toward four cell lines by MTT assay.

Experimental

Plant material

The aerial parts of *S. laxa* Boiss. and Buhse., from Siah-bishe in Mazandaran province, *S. trinervis* Aitch. and Hemsl from Karaj in Alborz province, and *S. subaphylla* Rech. F. and *S. turcomanica* Trautv. from Golestan province

were collected in May 2008. The plants have been identified and deposited at the Herbarium of Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran.

Extraction

Freshly collected aerial parts of four species of *Stachys* were cleaned and shade dried. These parts were coarse powdered in a hand mill and stored at room temperature. Two hundred grams of powdered plants were extracted through percolation method with 80% aq. MeOH three times at room temperature. The extract was evaporated using rotary evaporator and consequently partitioned between CHCl_3 , EtOAc and MeOH. Each fraction evaporated with rotary evaporator and has been stored at refrigerator for the investigation of cytotoxic activity.

Cytotoxicity assay

The colon carcinoma (HT-29), colorectal adenocarcinoma (Caco-2) and ductal carcinoma (T47D) cell lines were mentioned as exponentially growing cultures in RPMI 1640 cell culture medium (PAA, Germany), supplemented with 10% fetal bovine serum (FBS: Gibco, USA), for HT-29 cells and 15% FBS for Caco-2 and T47D cells. The Swiss mouse embryo fibroblast (NIH 3T3) cell line was kept in Dulbecco's modified Eagle's medium (DMEM; PAA, Germany) supplemented with 10% FBS. 100 IU/mL penicillin and 100 $\mu\text{g}/\text{mL}$ streptomycin (Roche, Germany) were added to the media. All the cell lines were cultured at 37°C in air /carbon dioxide (95:5) atmosphere.

Cytotoxic activity was measured using modified MTT assay (31). 1×10^4 cells/well were plated in 96-well plates (Nunc, Denmark) and incubated for 24 h before the addition of drugs. After 96 h of incubation in Caco₂ cells and 48 h of incubation in HT-29, NIH/3T3 and T47D cells, 20 μL of MTT (Merck, Germany) reagent (5 mg/mL) in phosphate buffered saline (PBS) was added to each well. The plates were incubated at 37°C for 4 h. The medium was discharged and the formazan blue, which had been formed in the cells, were dissolved with 100 μL dimethyl sulphoxide (DMSO). After the incubation at 37°C for 10 min, absorbance at 570 nm at the dissolved solutions was detected using a micro plate reader

Table 1. Cytotoxic activity of total extract and fractions of four species of *Stachys*.

Sample	Cell Lines* (MTT assay)			
	HT-29	Caco-2	T47D	NIH/ 3T3
<i>Stachys laxa</i>				
Total extract	421.97 ± 8.71	> 1000	239.78 ± 16.92	508.77 ± 45.56
Methanol fr.	265.83 ± 46.52	> 1000	254.1 ± 7.45	405.7 ± 74.18
Ethyl acetate fr.	134.004 ± 1.764	116.53 ± 18.23	18.079 ± 2.248	31.452 ± 1.554
Chloroform fr.	27.007 ± 2.096	101.65 ± 12.4	21.106 ± 2.491	41.294 ± 8.391
<i>Stachys subaphylla</i>				
Total extract	> 1000	> 1000	60.15 ± 3.72	771.26 ± 164.67
Methanol fr.	> 1000	> 1000	771.26 ± 197.38	> 1000
Ethyl acetate fr.	-	116.52 ± 2.78	51.05 ± 8.89	-
Chloroform fr.	234.86 ± 11.28	183.85 ± 8.87	43.411 ± 9.99	74.27 ± 2.34
<i>Stachys trinervis</i>				
Total extract	> 1000	> 1000	358.1 ± 14.14	> 1000
Methanol fr.	> 1000	> 1000	630.96 ± 29.99	649.23 ± 17.91
Ethyl acetate fr.	241.66 ± 14.71	338.22 ± 1.02	128.35 ± 6.65	110.05 ± 5.56
Chloroform fr.	> 1000	> 1000	383 ± 4.01	674.84 ± 67.37
<i>Stachys turcomanica</i>				
Total extract	219.58 ± 14.21	> 1000	103.67 ± 12.43	308.7 ± 1.34
Methanol fr.	693.57 ± 56.91	> 1000	708.60 ± 25.8	802.58 ± 26.84
Ethyl acetate fr.	66.10 ± 5.43	87.08 ± 8.9	30.14 ± 1.78	50.45 ± 3.45
Chloroform fr.	66.84 ± 7.92	187.89 ± 11.72	51.38 ± 9.49	58.22 ± 4.06
<i>Methotrexate</i>	0.23 ± 0.02	0.32 ± 0.04	0.16 ± 0.09	0.24 ± 0.013

*Results are expressed as IC₅₀ values (µg/mL). Key to cell Lines employed: HT-29 and Caco-2 (colon Adenocarcinoma), T47D (breast carcinoma), NIH 3T3 (Swiss embryo fibroblast).

(Anthos, Austria). The cell viability in MTT assay was calculated as the percentage of control value. Methotrexate was used as the positive control. Cytotoxicity was expressed as the concentration of extract inhibiting cell growth with 50% (IC₅₀ ± SD). All tests and analysis were run in triplicate.

Statistical analysis

IC₅₀ (the median growth inhibitory concentration) values were calculated from the IC₅₀ of dose-response curve in the sigma plot 11 software. Data representative of three independent experiments with similar results were presented as mean ± SD.

Result

The effects of these plant extracts on the proliferative response of the HT-29, Caco-2 and T47D cell lines have been analyzed by

treating the cells with different concentrations of the extracts and significant decrease in cell lines proliferation were observed. IC₅₀ ± SD are reported in Table 1. The chloroform and ethyl acetate fractions of *S. laxa* Boiss. showed high cytotoxicity on T47D, HT-29 (IC₅₀ < 50 µg/mL). The cytotoxicity of ethyl acetate fraction of *S. turcomanica* Trautv. and chloroform fraction of *S. subaphylla* were better than the other fractions on T47D cell line (IC₅₀ < 50 µg/mL). Total extract and fractions of *S. trinervis* did not affect the cell lines.

Discussion

Among all the samples, nonpolar (chloroform and ethyl acetate fractions) fractions of *S. laxa* exhibited greatest cytotoxicities on T47D and HT-29 cell lines compared with polar fraction and total extract. According to the data, the

cytotoxic activity of chloroform and ethyl acetate fractions on HT-29 and T47D cell lines were much stronger than that of Caco-2. It indicated that chloroform and ethyl acetate fractions of *S. laxa* had potential cytotoxic selectivity on T47D cell line. There was a report about the antioxidation and total phenol content of some *Stachys* spp. The research implied that total phenol content and FRAP value of methanolic extract are in this order: *S. laxa* > *S. turcomanica* > *S. subaphylla* > *S. trinervis* (33). Except *S. subaphylla* total extract against T47D cell line, the other methanolic extracts have indicated the same order of cytotoxic activity on T47D and HT-29. Higher cytotoxic activity of nonpolar fraction of *S. laxa* and *S. turcomanica* may be due to the high content of germacrene D in their essential oil, same as the *Stachys cretica* ssp. (34, 35), but main components of *S. subaphylla* and *S. trinervis* essential oils were identified as monoterpene hydrocarbons. In comparison with another fraction, methanolic and total fractions of all samples demonstrated slightly cytotoxic effect on cell line tested. The real IC₅₀ values of fractions of four species *Stachys* may be considerably lower than the positive control (Methotrexate) since its pharmacological active compounds are not pure and further researches are needed for defining potential component as cytotoxic natural medicines.

References

- (1) Rechinger KH and Hedge IC. *Flora Iranica*. Akademisch Druck-und Verlagsanstalt, Graz, Austria (1982) 150: 360-361.
- (2) Mozaffarian V. *a Dictionary of Iranian Plant Names*. Farhang Moaser, Tehran (1996) 522.
- (3) Vundac VB, Brantner AH and Plazibat M. Content of phenolic constituents and antioxidant activity of some *Stachys* taxa. *Food Chem.* (2007) 104: 1277-81.
- (4) Vundac VB, Males Z, Plazibat M, Golja P and Cetina-Cizmek BC. HPTLC determination of flavonoids and phenolic acids in some Croatian *Stachys* taxa. *J. Planar Chromatogr. Mod. TLC* (2005) 18: 269-73.
- (5) El-Ansari MA, Nawwar MA and Saleh NAM. Stachysetin, a diapigenine-7-glucoside-p-p-dihydroxytruxinate from *Stachys aegyptiaca*. *Phytochem.* (1995) 40: 1543-48.
- (6) Miyase T, Yamamoto R and Ueno A. Phenyl ethanoid glycosides from *Stachys officinalis*. *Phytochem.* (1996) 43: 475-79.
- (7) Nishimura H, Sasaki H, Inagaki N, Chin M and Mitsuhashi H. Nine phenethyl alcohol glycosides from *Stachys seiboldii*. *Phytochem.* (1991) 30: 659-69.
- (8) Amirghofran Z, Bahmani M, Azadmehr A and Javidnia K. Anticancer effects of various Iranian native medicinal plants on human tumor cell lines. *Neoplasma* (2006) 53: 428-33.
- (9) Amirghofran Z, Bahmani M, Azadmehr A and Javidnia K. Immunomodulatory and apoptotic effects of *Stachys obtusirena* on proliferative lymphocytes. *Med. Sci. Monit.* (2007) 13: 145-50.
- (10) Stamatias G, Kyriazopoulos P, Golegou S, Basayiannis A, Skaltsas S and Skaltsa H. *In-vitro* anti-*Helicobacter pylori* activity of Greek herbal medicines. *J. Ethnopharmacol.* (2003) 88: 175-9.
- (11) Grujic-Jovanovic S, Skaltsa HD, Marin P and Sokovic M. Composition and antibacterial activity of the essential oil of six *Stachys* species from Serbia. *Flav. Fragr. J.* (2004) 19: 139-44.
- (12) Sonboli A, Salehi P and Nejad Ebrahimi S. Essential oil composition and antibacterial activity of the leaves of *Stachys schtschegleevii* from Iran. *Chem. Nat. Compd.* (2005) 41: 171-4.
- (13) Digrak M, Hakki Alma M and Ilcim A. Antibacterial and antifungal activities of Turkish medicinal plants. *Pharm. Biol.* (2001) 39: 346-50.
- (14) Aydin A, Sener B, Cakici I, Turan NN and Erdemoglu N. Antioxidant activities of some Lamiaceae plant extracts. *Phytother. Res.* (2006) 20: 91-3.
- (15) Kukic J, Petrovic S and Niketic M. Antioxidant activity of four endemic *Stachys* taxa. *Biol. Pharm. Bull.* (2006) 29: 725-9.
- (16) Matkowski A and Piotrowska M. Antioxidant and free radical scavenging activities of some medicinal plants from the Lamiaceae. *Fitoterapia* (2006) 77: 346-53.
- (17) Khanavi M, Sharifzadeh M, Hadjiakhoondi A and Shafiee A. Phytochemical investigation and anti-inflammatory activity of aerial parts of *Stachys byzanthina* C. Koch. *J. Ethnopharmacol.* (2005) 97: 463-8.
- (18) Khanavi M, Sharifzadeh M, Hadjiakhoondi A and Shafiee A. Anti-inflammatory activity of aerial part of *Stachys byzanthina* C. Koch. *Iranian J. Pharm. Res.* (2004) Supp. 2: 55-56.
- (19) Kukic J, Dobric S and Petrovic S. Influence of some *Stachys* taxa on carrageenan-induced paw edema in rats. *Pharm. Biol.* (2007) 45: 560-3.
- (20) Maleki N, Garjani A, Nazemiyah H, Nilouroushan N, Eftekhari Sadat AT, Allameh Z and Hasannia N. potent anti-inflammatory activities of hydroalcoholic extract from aerial parts of *Stachys inflata* on rats. *J. Ethnopharmacol.* (2001) 75: 213-8.
- (21) Sharifzadeh M, Sharifzadeh K, Khanavi M, Hadjiakhoondi A and Shafiee A. Anti-inflammatory activity of aerial parts of *Stachys setifera* and *Stachys persica*. *Int. J. Pharmacol.* (2005) 1: 132-7.
- (22) Skaltsa HD, Bermejo P, Lazari DM, Silvan AM, Skaltsounis AL, Sanz A and Abad MJ. Inhibition of prostaglandin E₂ and leukotriene C₄ in mouse peritoneal macrophages and thromboxan B₂ production in human

- platelets by flavonoids from *Stachys chrysantha* and *Stachys candida*. *Biol. Pharm. Bull.* (2000) 23: 47-53.
- (23) Hayashi K, Nagamatsu T, Ito M, Hattori T and Suzuki Y. Acetoside, a component of *Stachys sieboldii* MIQ, may be a promising antinephritic agent. Effects of acetoside on crescentic-type anti-GBM nephritis in rats. *Jpn. J. Pharmacol.* (1994) 65: 143-51.
- (24) Rabbani M, Sajjadi SE and Zarei HR. Anxiolytic effects of *Stachys lavandulifolia* Vahl on the elevated plus-maze model of anxiety in mice. *J. Ethnopharmacol.* (2003) 89: 271-6.
- (25) Gruenwald J, Brendler T and Jaenicke C. *PDR for Herbal Medicines*. 2nd ed., Medical Economics Company, New Jersey (2000) 832.
- (26) Şerbetçi T, Demirci B, Güzelc ÇB, Kültür Ş, Ergüven M and Başer KHC. Essential oil composition, antimicrobial and cytotoxic activities of two endemic *Stachys cretica* subspecies (Lamiaceae) from Turkey. *Nat. Prod. Commun.* (2010) 5: 1369-74.
- (27) Firouznia A, Rustaiyan A, Masoudi S, Rahimizade M, Bigdeli M and Tabatabaei-Anaraki M. Volatile constituents of *Salvia limbata*, *Stachys turcomanica*, *Scutellaria litwinowii* and *Hymenocrater elegans* four Lamiaceae herbs from Iran. *J. Essent. Oil-Bearing Plants* (2009) 12: 482-9.
- (28) Khanavi M, Farahanikia B, Janbakhsh S, Sheibani S, Hoseini-Sajedi SM, Salahi-Oliaee MH, Ajani Y and Hadjiakhoondi A. Comparison of the essential oil composition of *Stachys trinervis* Aitch. & Hemsl. and *Stachys subaphylla* Rech. F. *J. Essent. Oil-Bearing Plants* (2008) 11: 406-12.
- (29) Sajjadi SE and Mehregan I. Composition of the essential oil of *Stachys laxa* Boiss. & Buhse. *Iranian J. Pharm. Res.* (2003) 8: 57-58.
- (30) Háznagy-Radnai E, Réthy B, Czigle Sz, Zupkó I, Wéber E, Martinek T, Falkay Gy and Máthé I. Cytotoxic activities of *Stachys* species. *Fitoterapia* (2008) 79: 595-7.
- (31) Newman DJ and Cragg GM. Natural products as sources of new drugs over the last 25 years. *J. Nat. Prod.* (2007) 70: 461-77.
- (32) Rahman A, Choudhary MI and Thomsen WJ. *Bioassay Techniques for Drug Development*. Taylor and Francis Group, Netherlands (2001) 34-35.
- (33) Khanavi M, Hajimahmoodi M, Cheraghi-Niroomand M, Kargar Z, Ajani Y, Hadjiakhoondi A and Oveisi MR. Comparison of the antioxidant activity and total phenolic contents in some *Stachys* species. *Afr. J. Biotechnol.* (2009) 8: 1143-7.
- (34) Kuźma Ł, Kalembe D, Różalski M, Różalska B, Więckowska-Szakiel M, Krajewska U and Wysokińska H. Chemical composition and biological activities of essential oil from *Salvia sclarea* plants regenerated *in-vitro*. *Molecules* (2009) 14: 1438-47.
- (35) Richmond JD, Agius BR, Wright BS, Haber WA, Moriarity DM and Setzer WN. Essential oil compositions and cytotoxic activities of *Dendropanax capillaris*, *Oreopanax nubigenus*, and *Schefflera rodrigueziana* from Monteverde, Costa Rica. *Nat. Prod. Commun.* (2007) 4: 271-4.