

Effects of *Siraitia grosvenorii* Fruits Extracts on Physical Fatigue in Mice

Da-Duo Liu^a, Xue-Wu Ji^{a*} and Rong-Wei Li^b

^aPhysical Education Institute of Jilin Normal University, Siping City, Jilin 136000, China.

^bDepartment of Physical Education of Hebei Normal University of Science and Technology, Qinhuangdao City, Hebei 066004, China.

Abstract

In this study, the effects of *Siraitia grosvenorii* fruits extracts (SGFE) on physical fatigue were investigated. One hundred and forty-four mice were randomly divided into four groups: control group, low-dose SGE-treated group, middle-dose SGE-treated group and high-dose SGFE-treated group. The animals of control group received an oral administration of physiological saline in a volume of 2.5 mL, and the animals of treated group received the same volume of SGFE (100, 200 and 400 mg/Kg bodyweight, once a day) for 28 days. After 28 days, anti-fatigue effects of SGFE were assessed 10 h after the last treatment by forced swimming test and some biochemical parameters related to fatigue, including blood lactic acid, serum urea nitrogen, liver glycogen and muscle glycogen were measured. The data showed that SGFE can extend the swimming time of the mice, as well as increasing the liver and muscle glycogen contents, but decrease the blood lactic acid and serum urea nitrogen levels. These results indicated that *Siraitia grosvenorii* fruits extracts had significant anti-fatigue effects on mice and these effects were dose-dependent.

Keywords: *Siraitia grosvenorii* fruits extracts; Physical fatigue; Mice; Forced swimming test.

Introduction

Fatigue is a common symptom both in sickness and in health and it is best defined as difficulty in initiating or sustaining voluntary activities (1, 2). It can be subdivided into physical and mental fatigues. For physical fatigue, it is generally accepted that muscle fatigue mainly arises in muscle tissue from a disorder of energy metabolism, not limited to lactic acid accumulation, which is caused by long-duration or intense exercise (3, 4). In addition to this muscle fatigue, the sensation of fatigue, which partly resembles mental fatigue in showing

lassitude, sleepiness and decreased motivation, often develops after exercise (5-7). Recently, there has been a great increase in the use of over-the-counter supplements and naturally occurring nutraceuticals for the attenuation of physical fatigue (8).

Siraitia grosvenorii (Swingle), a perennial vine of the Cucurbitaceae family, has been cultivated in the restricted area of the southern part of China (9, 10). Fruits of these plants, known as Luo-Han-Guo in China, have been widely used in the treatment of laryngitis, bronchitis, and gastrointestinal diseases in traditional Chinese medicine (11, 12). Studies showed that *Siraitia grosvenorii* fruits extracts (SGFE) possess a wide range of pharmacologic and health-promoting properties including immune

* Corresponding author:

E-mail: xuewuji@yahoo.cn

enhancement, anti-oxidation, anti-diabetes, anti-tumor and anti-inflammation (13, 14). The main bioactive components of this plant fruits are mogrosides, which are members of the family of triterpene glycosides, including mogroside IVa, mogrosides V and VI, iso-mogroside V, 11-oxomogroside V and siamenoside I (15). Previous studies showed that glycosides are bioactive components of many famous Chinese medicines in recent years and used in developing new drugs (16-19). Triterpene glycosides such as *Siraitia grosvenorii* have been used as an anti-hepatitis drug and steroidal glycosides have been used as anti-epilepsy, anti-inflammatory and anti-tumor glycosides (20-23). However, very little is known about anti-fatigue effects of SGFE. The present study aimed to investigate the effects of SGFE on physical fatigue in mice.

Experimental

Chemical

All chemicals were purchased from Jilin Chemical Reagents Co., Ltd. (Changchun, China) unless otherwise indicated. Commercial diagnostic kits (special for animal testing) used to determine blood lactic acid (BLA) and tissue glycogen were purchased from Nanjing Jiancheng Bioengineering Institute (Nanjing, China). Commercial diagnostic kit used to determine serum urea nitrogen (SUN) was purchased from Biosino Biotechnology Co., Ltd. (Beijing, China). Water was purified with a Milli-Q purification system and was used to prepare all solutions.

Plant materials

Fruits of *Siraitia grosvenorii* were purchased from Jilin Pharmaceutical Co. (Changchun, China), and identified by Professor Yang LY, College of Traditional Chinese Medicine, Jilin Agricultural University. Voucher specimens were deposited at the herbarium of Jilin Normal University.

Preparation of Siraitia grosvenorii fruits extracts

Siraitia grosvenorii fruits extracts (SGFE) were prepared as described previously (24) with slight modifications. Air-dried *Siraitia*

grosvenorii fruits (500 g) were ground into powder (particle diameter: 0.2 to 0.5 mm), then the powder was dissolved into 3000 mL distilled water and extracted 3 times at 70°C for 1 h every time. Subsequently, the water-soluble extracts were vacuum concentrated, separated and purified using column chromatography, at last vacuum-dried to yield extracts of *Siraitia grosvenorii* fruits (46.4 g/Kg), which were determined by HPLC system (Waters, Milford, MA). The HPLC system consisted of Waters 1525 binary HPLC pump, Waters 2487 dual λ absorbance detector and a XTerra RP18 column. HPLC analysis indicated that this extracts contained 76.4% glycosides and the major glycoside was mogrosides V (25.27%). Solution of aqueous extract was prepared in saline for the experiment.

Animals

Male ICR mice (18-22 g) were purchased from the Jilin Laboratory Animal Breeding and Research Center (Changchun, China). The mice were individually housed in a room maintained at $23 \pm 2^\circ\text{C}$ and $50 \pm 5\%$ humidity with a 12 h light-dark cycle. They were given free access to food and water throughout the experiments. The experiments were carried out in accordance with the China animal protection law and approved by Ethics Commission of Jilin Normal University.

Experimental design

After an adaptation period for a week, the 144 mice were randomly divided into four groups ($n = 36$ per group): control (C) group, low-dose SGFE-treated (LT) group, middle-dose SGFE-treated (MT) group and high-dose SGFE-treated (HT) group. The animals of control (C) group received an oral administration of physiological saline in a volume of 1.0 mL, and the animals of treated group received the same volume of SGFE (100, 200 and 400 mg/Kg bodyweight/day) for 28 days. The rationale for the selection of the doses was based on our previous experiments and some early literature. The doses of SGFE (100-400 mg/Kg bodyweight) were confirmed to be suitable and effective in the tested mice. The mice were made to swim for 15 min three times a week to accustom them to swimming.




Figure 1. Effect of *Siraitia grosvenorii* fruit extracts (SGFE) on swimming time-to-exhaustion of mice. Data were presented as means \pm SD (n = 12 per group). C: control, LT: low-dose SGFE-treated (100 mg/Kg bodyweight); MT: middle-dose SGFE-treated (200 mg/Kg bodyweight), HT: high-dose SGFE treated (400 mg/Kg bodyweight). *: $p < 0.05$ as compared with the control group

After 28 days, anti-fatigue effects of SGFE were assessed 10 h after the last administration by forced swimming test.

Forced swimming test

Twelve mice were taken out from each group to make forced swimming test. The apparatus used in this test was an acrylic plastic pool (90 \times 45 \times 45 cm) filled with water maintained at $25 \pm 2^\circ\text{C}$. The water in the acrylic plastic pool was 35 cm deep. Mice had a load attached (10% body weight) to its tail to reduce the time of swimming-to-exhaustion tests. Exhaustion was determined by observing loss of coordinated movements and failure to return to the surface within 10 s (25-27) and the swimming time was immediately recorded.

Analysis of blood biochemical parameters related to fatigue

Twelve mice were taken out from each group for blood biochemical parameters analyses. Mice were forced to swim for 30 min after weight loading (2% body weight) (25). After resting for 60 min, blood was collected from orbital sinus to examine the blood lactic acid (BLA) and serum urea nitrogen (SUN). Then, BLA and SUN contents were tested according

to the recommended procedures provided by the commercial diagnostic kit.

Analysis of tissue glycogen contents

Twelve mice were taken out from each group for tissue glycogen contents analyses. Mice were forced to swim for 90 min without loads. After resting for 60 min, the mice were killed to collect their livers and gastrocnemius muscles (28). Then, tissue glycogen contents were tested according to the recommended procedures provided by the commercial diagnostic kit.

Statistical analysis

All results are expressed as mean \pm SD for ten mice in each group. To determine the effect of treatment, data was analyzed using one-way ANOVA repeated measures. P-values of less than 0.05 were regarded as significant. Significant values were assessed with Duncan's multiple range test. Data was analyzed using the statistical package "SPSS 12.0 for Windows".

Results and Discussion

Effect of SGFE on body weights of mice

The body weights of mice were measured at the beginning of the experiment (day 0) and




Figure 2. Effect of *Siraitia grosvenorii* fruit extracts (SGFE) on blood lactic acid of mice.

Data were presented as means \pm SD (n = 12 per group). C: control, LT: low-dose SGFE-treated (100 mg/Kg bodyweight); MT: middle-dose SGFE-treated (200 mg/Kg bodyweight), HT: high-dose SGFE-treated (400 mg/Kg bodyweight). *: p < 0.05 as compared with the control group.

after the administration with different dosages of SGE for 28 days (28th day). It was found (data not shown) that the body weights of mice in the SGFE-treated groups (LT, MT and HT group) were not different significantly from that in the control group (C group) at day 0 or 28 (p > 0.05), which means the SGFE has no effect on body weight.

Effect of SGFE on the time of swimming-to-exhaustion tests swimming time to exhaustion of mice

A direct measure of an anti-fatigue effect is the increase in exercise tolerance. Swimming-to-exhaustion is an experimental exercise model to evaluate the anti-fatigue effect; it works well for evaluating the endurance capacity of mice and gives a high reproducibility (29). To standardize the workload and reduce the swimming time, weights at specific body weight percentages were added to the chest or tail of the animal (30). Reduced susceptibility to fatigue is correlated with longer swimming times (31). As shown in Figure 1, the swimming time-to-exhaustion of the SGFE-treated groups (LT, MT and HT group) were significantly longer than that of control group (C group) (p < 0.05). The results indicated that SGFE had anti-fatigue effects and these effects were dose-dependent.

Effect of SGFE on blood lactic acid and serum urea nitrogen of mice

The swimming exercise is known to induce the blood biochemical changes (32). Blood lactic acid (BLA) is the glycolysis product of carbohydrate under an anaerobic condition, and glycolysis is the main energy source for intense exercise in a short time. Therefore, BLA is one of the important indicators for judging the degree of fatigue (28, 33). In other words, BLA represents the degree of fatigue after exercise and the condition of recovery (26). As shown in Figure 2, after swimming, the SGFE-treated groups (LT, MT and HT group) showed a dose-dependent decrease in the BLA levels compared with the control group (C group) (p < 0.05). The results indicated that SGFE can effectively delay the increase of lactic acid in the blood and postpone the appearance of physical fatigue.

Blood urea nitrogen (BUN), the metabolism outcome of protein and amino acid, is a sensitive index to evaluate the bearing capability when human bodies suffer from a physical load. Several studies have shown that the BUN in the blood rises significantly for a long-run athlete after exercise (26, 34, 35). In other words, the worse the body is adapted for exercise tolerance, the more significantly the BUN level increases (29, 36). Therefore, BUN is another index of fatigue status. As shown in Figure 3, after swimming,




Figure 3. Effect of *Siraitia grosvenorii* fruit extracts (SGFE) on serum urea nitrogen (SUN) of mice. Data were presented as means \pm SD (n = 12 per group). C: control, LT: low-dose SGFE treated (100 mg/Kg bodyweight); MT: middle-dose SGFE treated (200 mg/Kg bodyweight), HT: high-dose SGFE treated (400 mg/Kg bodyweight). *: p < 0.05 as compared with control group

MT and HT group showed a significant decrease in the BUN levels compared with the control group (C group) (p < 0.05). However, BUN levels of the LT group were lower than that of the control group (C group), but there was no significant difference (p > 0.05). The results indicated that SGFE may reduce the catabolic decomposition of protein for energy.

Effect of SGFE on liver and muscle glycogen of mice

It was known that the endurance capacity of body was markedly decreased if the energy was exhausted (37). Energy for exercise is derived initially from the breakdown of glycogen, after the strenuous exercise muscle glycogen exhausts, and later, energy forms circulating glucose released by the liver (38, 39). Thus, the glycogen contents are sensitive parameters

related to fatigue. Table 1 shows the effect of SGFE on liver and muscle glycogen of mice. After swimming, the SGFE-treated groups (LT, MT and HT group) showed a dose-dependent increase in the liver and muscle glycogen contents compared with the control group (C group) (p < 0.05). The results indicated that the rate of glycogen depletion was delayed in the SGFE-treated groups, and suggested that SGFE could decrease carbohydrate utilization during the exercise.

In conclusion, the data showed that SGFE can extend the swimming time of the mice, as well as increasing the liver and muscle glycogen contents, but it decreases the blood lactic acid and serum urea nitrogen levels. These results indicated that SGFE had significant anti-fatigue effects on mice and these effects were dose-dependent. However, further study is needed

Table 1. Effect of *Siraitia grosvenorii* fruits extracts (SGFE) on liver and muscle glycogen of mice.

Group	Liver glycogen (mg/g)	Muscle glycogen (mg/g)
C	7.54 \pm 2.21	1.21 \pm 0.27
LT	14.27 \pm 3.38*	1.86 \pm 0.42*
MT	17.29 \pm 3.87*	1.95 \pm 0.33*
HT	19.63 \pm 3.11*	2.06 \pm 0.61*

Data were presented as means \pm SD (n = 12 per group). C: control, LT: low-dose SGFE treated (100 mg/Kg bodyweight); MT: middle-dose SGFE treated (200 mg/Kg bodyweight), HT: high-dose SGFE treated (400 mg/Kg bodyweight). *: p < 0.05 as compared with the control group.

to elucidate the more exact mechanism of the anti-fatigue effect of SGFE at the cellular and molecular levels.

References

- (1) Sugino T, Shirai T, Kajimoto Y and Kajimoto O. L-ornithine supplementation attenuates physical fatigue in healthy volunteers by modulating lipid and amino acid metabolism. *Nutr. Res.* (2008) 28: 738-43.
- (2) Tanaka M, Baba Y, Kataoka Y, Kinbara N, Sagesaka YM, Kakuda T and Watanabe Y. Effects of (-)-epigallocatechin gallate in liver of an animal model of combined (physical and mental) fatigue. *Nutrition.* (2008) 24: 599-603.
- (3) Powers SK and Jackson MJ. Exercise-induced oxidative stress: cellular mechanisms and impact on muscle force production. *Physiol. Rev.* (2008) 88: 1243-76.
- (4) Allen DG, Lamb GD and Westerblad H. Skeletal muscle fatigue: cellular mechanisms. *Physiol. Rev.* (2008) 88: 287-332.
- (5) Morimoto Y, Zhang Q and Adachi K. Increased Blood Quinolinic Acid after Exercise in Mice: Implications for Sensation of Fatigue after Exercise. *J. Health Sci.* (2011) 57: 367-71.
- (6) Powers SK, Ji LL and Leeuwenburgh C. Exercise training-induced alterations in skeletal muscle antioxidant capacity: a brief review. *Med. Sci. Sports Exerc.* (1999) 31: 987-97.
- (7) Reid MB. Free radicals and muscle fatigue: Of ROS, canaries, and the IOC. *Free Radic. Biol. Med.* (2008) 44: 169-79.
- (8) Sugino T, Aoyagi S, Shirai T, Kajimoto Y and Kajimoto O. Effects of Citric Acid and L-Carnitine on Physical Fatigue. *J. Clin. Biochem. Nutr.* (2007) 41: 224-30.
- (9) Zhang L, Hu TJ and Lu CN. Immunomodulatory and antioxidant activity of a *Siraitia grosvenorii* polysaccharide in mice. *Afr. J. Biotech.* (2011) 10: 10045-53.
- (10) Qi XY, Chen WJ, Zhang LQ and Xie BJ. Mogrosides extract from *Siraitia grosvenorii* scavenges free radicals *in-vitro* and lowers oxidative stress, serum glucose, and lipid levels in alloxan-induced diabetic mice. *Nutr. Res.* (2008) 28: 278-84.
- (11) Hossen M A, Shinmei Y, Jiang S, Takubo M, Tsumuro T, Murata Y, Sugiura M and Kamei C. Effect of Lo Han Kuo (*Siraitia grosvenorii* Swingle) on nasal rubbing and scratching behavior in ICR mice. *Biol. Pharm. Bull.* (2005) 28: 238-41.
- (12) Matsumoto S, Jin M, Dewa Y, Nishimura J, Moto M, Murata Y, Shibutani M and Mitsumori K. Suppressive effect of *Siraitia grosvenorii* extract on dicyclanil-promoted hepatocellular proliferative lesions in male mice. *J. Toxicol. Sci.* (2009) 34: 109-18.
- (13) Lin GP, Jiang T, Hu XB, Qiao XH and Tuo QH. Effect of *Siraitia grosvenorii* polysaccharide on glucose and lipid of diabetic rabbits induced by feeding high fat/high sucrose chow. *Exp. Diabetes Res.* (2007) 67435: 1-4.
- (14) Akihisa T, Hayakawa Y, Tokuda H, Banno N, Shimizu N, Suzuki T and Kimura Y. Cucurbitane glycosides from the fruits of *Siraitia grosvenorii* and their inhibitory effects on Epstein-Barr virus activation. *J. Nat. Prod.* (2007) 70: 783-8.
- (15) Chaturvedula VSP and Prakash I. Kaempferol glycosides from *Siraitia grosvenorii*. *J. Chem. Pharm. Res.* (2011) 3: 799-804.
- (16) Hao M, Ba Q, Yin J, Li J, Zhao Y and Wang H. Deglycosylated ginsenosides are more potent inducers of CYP1A1, CYP1A2 and CYP3A4 expression in HepG2 cells than glycosylated ginsenosides. *Drug Metab. Pharmacokinet.* (2011) 26: 201-5.
- (17) Chen L, Qi J, Chang YX, Zhu D and Yu B. Identification and determination of the major constituents in Traditional Chinese Medicinal formula Danggui-Shaoyao-San by HPLC-DAD-ESI-MS/MS. *J. Pharm. Biomed. Anal.* (2009) 50: 127-37.
- (18) Zhao W, Ye Q, Tan X, Jiang H, Li X, Chen K and Kinghorn AD. Three new sesquiterpene glycosides from *Dendrobium nobile* with immunomodulatory activity. *J. Nat. Prod.* (2001) 64: 1196-200.
- (19) Lechtenberg M, Schepmann D, Niehues M, Hellenbrand N, Wünsch B and Hensel A. Quality and functionality of saffron: quality control, species assortment and affinity of extract and isolated saffron compounds to NMDA and sigma1 (sigma-1) receptors. *Planta. Med.* (2008) 74: 764-72.
- (20) Zhou J. Bioactive glycosides from Chinese medicines. *Mem. Inst. Oswaldo. Cruz.* (1991) 2: 231-4.
- (21) Chen G, Xu N, Li ZF, Zhang QH, Wu HH and Pei YH. Steroidal glycosides with anti-tumor activity from the roots of *Cynanchum wallichii* Wight. *J. Asian Nat. Prod. Res.* (2010) 12: 453-7.
- (22) Kawabata T, Cui MY, Hasegawa T, Takano F and Ohta T. Anti-inflammatory and anti-melanogenic steroidal saponin glycosides from Fenugreek (*Trigonella foenum-graecum* L.) seeds. *Planta. Med.* (2011) 77: 705-10.
- (23) Li RW, Leach DN, Myers SP, Lin GD, Leach GJ and Waterman PG. A new anti-inflammatory glucoside from *Ficus racemosa* L. *Planta. Med.* (2004) 70: 421-6.
- (24) Song F, Qi X, Chen W, Jia W, Yao P, Nussler AK, Sun X and Liu L. Effect of *Momordica grosvenorii* on oxidative stress pathways in renal mitochondria of normal and alloxan-induced diabetic mice. Involvement of heme oxygenase-1. *Eu.r J. Nutr.* (2007) 46: 61-9.
- (25) Yao LQ and Li FL. Lycium barbarum polysaccharides ameliorates physical fatigue. *Afr. J. Agric. Res.* (2010) 5: 2153-7.
- (26) Wang JJ, Shieh MJ, Kuo SL, Lee CL and Pan TM. Effect of red mold rice on antifatigue and exercise-related changes in lipid peroxidation in endurance exercise. *Appl. Microbiol. Biotechnol.* (2006) 70: 247-253.
- (27) Zhang XL, Ren F, Huang W, Ding RT, Zhou QS and Liu XW. Anti-Fatigue Activity of Extracts of Stem

- Bark from *Acanthopanax senticosus*. *Molecules*. (2011) 16: 28-37.
- (28) Ma L, Ca DL, Li HX, Tong BD, Song LH and Wang Y. Anti-fatigue effects of salidroside in mice. *J. Med. Coll. PLA*. (2008) 23: 88-93.
- (29) Zhang Y, Yao X, Bao B and Zhang Y. Anti-fatigue activity of a triterpenoid-rich extract from Chinese bamboo shavings (*Caulis bambusae in taeniam*). *Phytother. Res*. (2006) 20: 872-876.
- (30) Mizunoya W, Oyaizu S, Ishihara K and Fushiki T. Protocol for measuring the endurance capacity of mice in an adjustable-current swimming pool. *Biosci. Biotechnol. Biochem*. (2002) 66: 1133-1136.
- (31) You LJ, Zhao MM, Regenstein JM and Ren JY. *In-vitro* antioxidant activity and *in-vivo* anti-fatigue effect of loach (*Misgurnus anguillicaudatus*) peptides prepared by papain digestion. *Food Chem*. (2011) 124: 188-194.
- (32) Yalcin O, Bor-Kucukatay M, Senturk UK and Baskurt OK. Effects of swimming exercise on red blood cell rheology in trained and untrained rats. *J. Appl Physiol*. (2000) 88: 2074-2080.
- (33) Ding JF, Li YY, Xu JJ, Su XR, Gao X and Yue FP. Study on effect of jellyfish collagen hydrolysate on anti-fatigue and anti-oxidation. *Food Hydrocol*. (2011) 25: 1350-1353.
- (34) Wu IT. The effects of serum biochemical value with different beverage to replenish and intermittent exercise in high intensity. *Tahan. J. College Engr. Business J*. (1999) 13: 387-400.
- (35) Cooke M, Iosia M, Buford T, Shelmadine B, Hudson G, Kerksick C, Rasmussen C, Greenwood M, Leutholtz B, Willoughby D and Kreider R. Effects of acute and 14-day coenzyme Q10 supplementation on exercise performance in both trained and untrained individuals. *J. Int. Soc. Sports Nutr*. (2008) 5: 8.
- (36) Huang LZ, Huang BK, Ye Q and Qin LP. Bioactivity-guided fractionation for anti-fatigue property of *Acanthopanax senticosus*. *J. Ethnopharmacol*. (2011) 133: 213-9.
- (37) Tang KJ, Nie RX, Jing LJ and Chen QS. Anti-athletic fatigue activity of saponins (Ginsenosides) from American ginseng (*Panax quinquefolium L.*). *Afr. J. Pharm*. (2009) 3: 301-306.
- (38) Suh S H, Paik I Y and Jacobs K. Regulation of blood glucose homeostasis during prolonged exercise. *Mol. Cells*. (2007) 23: 272-9.
- (39) Wei W, Zheng LY, Yu MY, Jiang N, Yang ZR and Luo X. Anti-fatigue activity of extract from the submerged fermentation of *Ganoderma Lucidum* using *Radix astragali* as substrate. *J. Anim. Plant Sci*, (2010) 6: 677-84.

This article is available online at <http://www.ijpr.ir>
