

Improvement of Capillary Electrophoretic Enantioseparation of Fluoxetine by a Cationic Additive

Farin Sattary Javid^{a,b}, Alireza Shafaati^{a,c*} and Afshin Zarghi^{a,c}

^aPharmaceutical Chemistry Department, School of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran. ^bStudents Research Committee, School of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran. ^cPharmaceutical Sciences Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Abstract

One of the problems encountered in CE separations of basic compounds is the adsorption of analytes in negatively charged capillary wall which could lead to poor repeatability of migration time and peak area. Additionally, separation of enantiomers of chiral of basic drugs is commonly carried out in low pH buffer which contributes to strong ionic interaction of the cationic drug ions with negatively charged chiral selectors. The two phenomena results in poor enantioseparations. To overcome the problems associated with chiral separations of basic drugs by CE, the effect of guanidine (GU) on the improvement of chiral separation of a model basic drug, fluoxetine (FLX), was investigated. In the present study, GU was used as a cationic additive to the running buffer containing a chiral selector, sulfated beta cyclodextrine. Better results obtained with GU as the buffer additive in enantioseparation of FLX.

Keywords: Capillary electrophoresis; Chiral separation; Fluoxetine; Guanidine.

Introduction

More than 30% of drugs in market are chiral (1) and mostly two enantiomers of a chiral drug show different pharmacological, toxicological or pharmacokinetic properties (2).

In addition, separation methods are required for quality control of single-enantiomer pharmaceutical raw materials and products during their production, storage and application (3). Thus, different analytical techniques have been developed for controlling synthesis, enantiomer purity check and pharmacodynamic studies of chiral drugs.

Capillary electrophoresis (CE) as a powerful technique, especially for analytical

enantioseparations offers high efficiency and high flexibility with regard to the separation conditions, as well as the low consumption of chemicals and solvents. Consequently, it is shown that the major application of the technique is chiral separations (4). However, CE has some limitations, especially for chiral and non-chiral resolution of basic drugs.

Basic analytes are positively charged over a wide range of pH (especially in acidic solutions) and electrostatically attracted to the negatively charged capillary walls. The irreproducible adsorption may cause variations to migration time and peak area (5), and consequently results in lower precision. Furthermore, impurity peaks may be masked by the peak tailing or resolution of enantiomers, in which two analytes migrating closely after each other may deteriorate. Additionally, analyte absorption could raise

* Corresponding author:

E-mail: ashafaati@yahoo.com

the limit of detection (LOD) and limit of quantification (LOQ) of analysis.

Different approaches have been applied to avoid undesirable interactions of the analytes with the capillary wall, including the use of high salt concentration, extremes of pH, buffer additives, and coated capillaries (5).

In the challenge of developing a separation method by capillary electrophoresis (CE), a variety of additives may be added to the background electrolyte to improve the separation (6) the most used additives are nonionic additive namely organic modifiers (7-8) and urea (9-10). The highly ionic triethylamine could optimize the resolution of separation of peptides (11). Our previous work (12) has shown that the addition of guanidine (GU) to the running buffer containing sulfated- β -cyclodextrin (SB-CD) as a chiral selector increased the resolution of enantiomers of basic chiral compounds. In the course of our ongoing interest in chiral separations, the present study was conducted to investigate the influence of GU on the SB-CD mediated enantioseparations of fluoxetine.

Fluoxetine (FLX), as an antidepressant, is a selective serotonin reuptake inhibitor which is less sedating compared to tricyclic antidepressants and has fewer antimuscarinic and cardiotoxic effects (13). FLX (Figure 1) is a basic drug which contains tertiary amine nitrogen (pK_a of 10.3) (14) that is positively charged in neutral and lower pHs. This drug includes one chiral center within its structures. (*R*)-fluoxetine and (*S*)-fluoxetine are similarly effective at blocking serotonin reuptake. As these enantiomers are metabolized differently, the use of *R*-enantiomer was expected to result in less variable plasma levels of fluoxetine and its active metabolites compared to that observed with racemic fluoxetine. Additionally, it is shown that (*R*)-fluoxetine and its metabolites inhibit CYP2D6, a cytochrome P450 system enzyme, to a lesser extent than (*S*)-fluoxetine and its metabolites (15-16).

Few reports on the application of CE methods were found in the literature for separation of FLX enantiomers (17-18). These methods presented the enantioseparation in acidic condition using combined neutral and negatively charged cyclodextrins.




Figure 1. Chemical structure of fluoxetine HCl.

Experimental

Materials

All solutions were prepared in Nanopure 18M Ω ultrapure water (Barnstead, Chicago, IL, USA). Sodium phosphate buffer with pH range of 2-4 was prepared from sodium monohydrogen phosphate purchased from Merck (Darmstadt, Germany), and adjusting the pH with orthophosphoric acid purchased from the same company, sulfated β -cyclodextrin (Sigma, Louis, MO, USA), were used as received. GU HCl was obtained from Merck (Darmstadt, Germany), Racemic FLX (as HCl) was kindly gifted by Noor Research and Education Institute (Tehran, Iran).

CE system

All experiments were performed on a Prince-C650 CE system equipped with UV/visible detector BISCHOFF (λ 1010). CE analysis was performed using untreated fused-silica capillary of 50 μ m internal diameter and 75-cm total length (25-cm effective length). The capillary was conditioned prior to the first use by rinsing with 1 M NaOH for 15 min, followed by 0.1 M NaOH and water for 5 min each. Between runs, the capillary was flushed for 3 min with running buffer to guarantee the good reproducibility. The electrophoretic integration was performed by Dax Data Acquisition Analysis software (version 8.0).

Capillary electrophoretic conditions

The applied voltage was - 15 kV at 25°C, and the migrant species were detected at 230 nm.

The running buffer was a 50 mM phosphate buffer adjusted at pH = 2-4 by 1 M sodium hydroxide. The running buffer contained 1-4%




Figure 2. Separation of FLX enantiomers by CE. Experimental conditions: running buffer phosphate 25 mM (pH = 2.5) containing 3% SB-CD; wavelength, 230 nm; voltage, -15 kV; sample FLX 0.1 mg/mL, other condition as detailed in experimental section.

sulfated beta-cyclodextrine as chiral selector and guanidine (40-100 mM) as buffer additive, which was freshly prepared daily and filtered through a 0.45 μm filter membrane.

Sample injection was performed hydrodynamically by applying 50 mbar of pressure for 12 s.

Sample preparations

Standard stock solutions of racemic fluoxetine containing 1 mg/mL were prepared by dissolving standard powder in water. Working solutions were prepared by diluting stock solution of fluoxetine with water.

Results and Discussion

Optimization of the method

Concentration of the selector

Based on our previous works on enantioseparation of basic drugs with CE (19-20), the initial attempt was based on applying a reversed-mode CE with the pH adjusted to 2.5 using a 25 mM phosphate buffer as the running buffer. The SB-CD concentration was studied over a range of 1-4% with maximum resolution achieved at 3% of SB-CD (see Figure 2). Lower concentrations of the selector resulted in very broad peak, whilst higher concentrations

of the selector led to the deterioration of the enantioseparation.

Concentration of the cationic additive

To improve chiral separation of the two enantiomers of FLX, different concentration of GU over a range of 20-100 mM was added to the running buffer. As shown in Figure 3, baseline separation of the enantiomers was obtained at any GU concentration. At higher GU concentrations, better peak shapes were observed. But to avoid Joule heating, GU concentration was adjusted to 80 mM which resulted in a reasonable separation and peak shapes, as well as minimum current and Joule heating.

Running buffer pH and concentration

Buffer pH is one of the most important factors in CE as it may affect the charge of the analytes and the chiral selector, and thus the binding characteristics (21). In order to obtain the optimum results, the buffer pH and concentration was reconsidered. Figure 4 shows the effect of buffer pH on the enantioseparation of FLX, with SB-CD and GU concentration kept constant at their best. Baseline resolution of the two peaks with acceptable peak shapes was observed at pH of 3. At this pH, the electroosmotic flow was low enough to allow applying carrier-mode




Figure 3. The effect of GU concentration on resolution of the two enantiomers of FLX Experimental conditions as described in Figure 2.

separation, whilst ionization of FLX was high enough to render the formation of complex with negatively charged SB-CD molecules.

Despite better peak shapes obtained with higher buffer concentrations, to avoid Joule heating, buffer concentration was kept at 25 mM. Besides, GU was used as hydrochloride salt which in turn contributed to the buffer strength.

Optimized resolution

Figure 5 shows the final electropherogram

obtained at the optimum conditions, *i.e.* 80 mM GU (as HCl) and 3% SB-CD in 25 mM phosphate buffer at pH of 3.0 with applied voltage of - 15 kV.

GU Effects on enantioseparation

The mechanism of chiral separation using CDs is based on the different inclusion of the hydrophobic part of the enantiomers into the CD cavity. The stereoselectivity is enhanced by secondary interactions between the functional




Figure 4. The effect of buffer pH on resolution of the two enantiomers of FLX Experimental conditions as described in Figure 2 (GU concentration was 80 mM).




Figure 5. Resolution of the FLX enantiomers at optimum conditions

groups of the analyte with the hydroxyl (or derivetized hydroxyl) groups on the outside rim of the CD (22). As shown in Figure 2, in the absence of GU, complete resolution of two enantiomers of FLX was not achieved which was attributed to the strong interaction of FLX cations with the sulfate groups on the CD molecule.

GU is a basic compound which bears positive charge in acidic solutions. The positive charge on the molecule is dispersed and delocalized (24). Thus, it is expected that the cation interacts with silanol groups on the inner surface of the bared capillary and prevents FLX cations to adsorption on the capillary wall, which contribute to the interaction of the drug and CD. In addition, it is suggested that GU modifies interaction between FLX cations and negatively charged SB-CD ions, which renders the inclusion of drug into the CD cavity (12).

Conclusion

It was shown that GU greatly influenced resolution of the peaks relevant to the enantiomers of FLX when a reversed-mode CE method was applied using SB-CD in acidic buffer as a chiral selector. The GU effect was contributed to the modification of the interaction of FLX cations

with the silanol groups on the capillary inner wall and with the negatively charged sulfate moieties on the CD molecules.

Acknowledgment

We wish to thank the Research Deputy of School of Pharmacy, Shahid Beheshti University of Medical Sciences (SBMU) for the financial support of this work. The authors also appreciate Noor Research and Education Institute for kind gift of racemic-FLX.

References

- (1) Gubitz G and Schmid MG. Recent progress in chiral separation principles in capillary electrophoresis. *Electrophoresis* (2000) 21: 4112-4135.
- (2) Fanali S. Controlling enantioselectivity in chiral capillary electrophoresis with inclusion-complexation. *J. Chromatogr. A* (1997) 792: 227-267.
- (3) Chankvetadze B. Enantioseparations by using capillary electrophoretic techniques. The story of 20 and a few more years. *J. Chromatogr. A* (2007) 1168: 45-70.
- (4) Scriba GKE. Pharmaceutical and biomedical applications of chiral capillary electrophoresis and capillary electrochromatography: An update. *Electrophoresis* (2003) 24: 2409-2421.
- (5) Altria KD. Enhanced pharmaceutical analysis by CE using dynamic surface coating system. *J. Pharmaceut. Biomed.* (2003) 31: 447-453.

- (6) Marina ML, Rios A and Valcarcel M. *Chapter 1, Fundamental of capillary electrophoresis*, In: Marina ML, Rios A and Valcarcel M. (Eds) *Comprehensive Analytical Chemistry, Volume XLV*, Amsterdam Elsevier (2005) 1-29.
- (7) EeckhautAV, DetaevernierMR, CrommenJ and Michotte Y. Influence of methanol on the enantioresolution of antihistamine with carboxymethyl- β - cyclodextrin in capillary electrophoresis. *Electrophoresis* (2004) 25: 2838-2847.
- (8) Bean SR, Lookhart GL and Bietz JA. Acetonitrile as a buffer additive for free zone capillary electrophoresis separation and characterization of maize (*Zea mays* L.) and sorghum (*Sorghum bicolor* L. (Moench.)) storage proteins. *J. Agric. Food Chem.* (2000) 48: 318-327.
- (9) Hammitzsch-Wiedemann M and Scriba GKE. Influence of buffer substances and urea on the β -cyclodextrin-mediated chiral separation of dipeptides in CE. *Electrophoresis* (2007) 28: 2619-2628.
- (10) García-Ruiz C, Crego AL and Marina ML. Comparison of charged cyclodextrin derivatives for the chiral separation of atropisomeric polychlorinated biphenyls by capillary electrophoresis. *Electrophoresis* (2003) 24: 2657-2664.
- (11) Noumie Surugau L, Bergström EDT and Goodall DM. Effects of organic solvent and cationic additive on capillary electrophoresis of peptides. *Malaysian J. Anal. Sci.* (2008) 3: 520-527.
- (12) Sattarijavid F, Shafaati A and Zarghi A. Application of guanidine to improve enantioseparation of a model basic drug, cetirizine, by CZE using sulfated β -cyclodextrin. *Electrophoresis Submitted for publication* (2012)
- (13) Sweetman SC. (ed.) *Martindale: the Complete Drug Reference*. 35th ed. Pharmaceutical Press, London (2007) 351-352.
- (14) Lenkey N, Karoly RP, Szasz BK, Vizi ES and Mike A. the mechanism of activity-dependent sodium channel inhibition by the antidepressants fluoxetine and desipramine. *Mol. Pharmacol.* (2006) 70: 2052-2063.
- (15) McConathy J and Owens MJ. Stereochemistry in drug action primary care companion. *J. Clin. Psychiatry* (2003) 5: 70-73.
- (16) Patel BK and Hut AJ. Stereoselectivity in drug action and disposition: an overview. In: Reddy IK and Mehvar R. (eds.) *Chirality in Drug Design and Development*. Marcel Dekker, New York (2004) pages 127-174.
- (17) Desiderio C, Rudaz S, Raggi MA and Fanali S. Enantiomeric separation of fluoxetine and norfluoxetine in plasma and serum samples with high detection sensitivity capillary electrophoresis. *Electrophoresis* (1999) 17: 3432-3438.
- (18) Inoue T and Chang JP. Chiral separation of fluoxetine and its analogs with charged cyclodextrins by capillary electrophoresis. *J. Liq. Chromatogr.* (2003) 26: 2351-2367.
- (19) Zandkarimi M, Shafaati A, Foroutan SM and Lucy CA. Improvement of electrophoretic enantioseparation of amlodipine by polybrene. *Iranian J. Pharm. Res.* (2012) 11: 129-136.
- (20) Zandkarimi M, Shafaati A, Foroutan SM and Lucy CA. Rapid enantioseparation of amlodipine by highly sulfated cyclodextrins using short-end injection capillary electrophoresis. *DARU* (2009) 17: 269-276.
- (21) Rawjee YY, Staerk DU and Vigh G. Capillary electrophoretic chiral separations with cyclodextrin additives: I. acids: Chiral selectivity as a function of pH and the concentration of β -cyclodextrin for fenoprofen and ibuprofen. *J. Chromatogr. A* (1993) 635: 291-306.
- (22) Guttman A. Capillary electrophoresis separation of enantiomers by cyclodextrin array chiral analysis. In: Landers JP. (ed.) *Handbook of Capillary Electrophoresis*. 2nd ed. CRC Press, Boca Raton, Florida (1996) 75-100.