

Effects of Aqueous Extract of *Berberis integerrima* Root on Some Physiological Parameters in Streptozotocin-Induced Diabetic Rats

Hossein Ashraf*, Reza Heidari, Vahid Nejati and Minoos Ilkhanipoor

Department of Biology, Faculty of Sciences, Urmia University, Urmia, Iran.

Abstract

Diabetes mellitus is a common endocrine disorder. Anti-diabetic agents from natural and synthetic sources are available for the treatment of this disease. *Berberis integerrima* is a medicinal shrub used in conventional therapy for a number of diseases. The aim of the present study was to investigate the effects of aqueous extract of *Berberis integerrima* root (AEBI) on some physiological parameters in normal and streptozotocin-induced (STZ-induced) diabetic male Wistar rats. STZ-induced diabetic rats showed significant increases in the levels of blood glucose, triglycerides (TG), total cholesterol (TC), low density lipoprotein LDL-cholesterol (LDL-C), creatinine (Cr), urea, alanine aminotransferase (ALT), aspartate aminotransferase (AST), alkaline phosphatase (ALP), total bilirubin while body weight, high density lipoprotein HDL-cholesterol (HDL-C) and total protein levels were significantly decreased compared to normal rats. Treatment of diabetic rats with different doses of aqueous extract of *Berberis integerrima* root (250 and 500 mg/Kg bw) resulted in a significant decrease in blood glucose, triglycerides, cholesterol, LDL-cholesterol, ALT, AST, ALP, total bilirubin, creatinine and urea while HDL-cholesterol and total protein levels were markedly increased after six weeks compared to untreated diabetic rats. The effects of the AEBI at dose of 500 mg/Kg in all parameters except blood glucose (similar) is more than to the standard drug, glibenclamide (0.6 mg/Kg, p.o.). The results of this study indicate that the tested aqueous extract of *Berberis integerrima* root possesses hypoglycemic, hypolipidemic and antioxidant effects in STZ-induced diabetic rats.

Keywords: *Berberis Integerrima*; Streptozotocin; Hypoglycemic; Hypolipidemic; Antioxidant.

Introduction

Diabetes Mellitus is the heterogeneous metabolic disorder characterized by altered carbohydrate, lipid and protein metabolism (1). More than 170 million people worldwide have diabetes and this number is set to be increase up to 360 million by 2030 (2). In Iran, approximately 2 million adult people have been diagnosed with

diabetes and about 4.4 million of them have impaired fasting glucose (3). Although the main and effective treatment for diabetes mellitus is currently using insulin and hypoglycemic factors, these compounds have undesirable effects such as increased fat deposits, loss of fat tissue at the injection site and causing a hypoglycemic shock and having no effects on the trends of debilitating complication of diabetes in the long-term. Given the increase in human knowledge about the heterogeneity of this illness, there is a need to find effective combinations with fewer

* Corresponding author:

E-mail: hossein.ashraf@gmail.com

side effects in the treatment of diabetes (4). In addition, there are several forms of dyslipidemia in patients with diabetic mellitus. Due to the heart-vascular risks induced by hyperglycemia and hyperlipidemia, lipid disorders should be quickly diagnosed and treated as a part of diabetes comprehensive treatment. The most common pattern of dyslipidemia is increased triglycerides and decreased HDL cholesterol (5). Although medicinal plants and their ingredients have been considered a long time in the treatment of diabetes mellitus, there are not reliable and valid studies on their absolute effectiveness (6). *Berberis integerrima* (Berberidaceae) is an important medicinal shrub, to a height of about 4 m, with thick and leathery leaves, a cluster inflorescence with 2-5 cm length and berry fruit with 7-8 mm length. This plant grows in most regions of Iran, especially in northern and northeast regions of the country. The harvest time is in November. Due to having secondary metabolites such as Berberine, Oxyacanthine, Berbamine, Palmatine, Jatrorrhizine, Columbamine and Berberubine, this plant has much medicinal properties (7) and is used in the treatment of gastrointestinal disease, bleeding, swollen gums, teeth, sore throat, fever, bile, malaria, hepatitis, inflammation and diarrhea. In addition, it has an important role in reducing blood cholesterol (7-10). Berberine is demonstrated to reduce serum cholesterol, triglyceride and LDL-cin subjects with dyslipidemia in animals (52).

Since few studies are available on anti-diabetic properties of this plant (*B. integerrima*), we examined the effects of aqueous extracts of *Berberis integerrima* root on some physiological parameters in streptozotocin-induced diabetic rats and compared these effects with sulfonylurea drugs like glibenclamide in a period of 6 weeks.

Experimental

Streptozotocin (STZ) was purchased from Sigma Chemical Co. (St. Louis, MO, USA). All the remaining chemicals were of highest commercially available grades.

Plant material

Wild samples of Barberry root (*Berberis integerrima*) were collected from suburb

Bavanat City (Fars Province, Iran) during November and December 2011 and identified by the Botany Department of Urmia University. A voucher specimen of the plant was deposited in the herbarium of the Faculty of Sciences, Urmia University, Urmia, Iran (No. 9059).

Preparation of aqueous extract

Roots were dried in the shade after being washed with cold water and were then powdered by using dry grinder and passed through the sieve. The aqueous extract was prepared by cold maceration of 150 g of powdered root barks in 500 mL of distilled water for 72 h. Then, the extract was filtered, concentrated, dried *in-vacuo* (yield 10 g) and the residue was stored in a refrigerator at 2-8°C for use in subsequent experiments (11).

Animals

Male Wistar rats weighing approximately 180-220 g (obtained from the central animal house of the Tehran Pasteur Institute, Tehran, Iran) were housed in an air-conditioned room under a 12-h light-dark cycle. Animals were allowed free access to tap water and standard laboratory rat food. All experimental procedures involving animals were approved by the Animal Research Ethics Committee of Urmia University of Faculty of Sciences, Urmia, Iran.

Acute toxicity study

Acute toxicity study of aqueous extract of *Berberis integerrima* root was determined as per the OECD guideline No. 423 (Acute Toxic Class Method). It was observed that test extract was not lethal to the rats even at 2500 mg/Kg dose. Hence, 1/10th (250 mg/Kg) and 1/5th (500 mg/Kg) of this dose were selected for further study (12).

Experimental induction of diabetes

Diabetes was induced in rats by intraperitoneal (IP) injection of streptozotocin (STZ) at a dose of 65 mg/Kg bw, dissolved in 0.1 cold citrate buffer (pH = 4.5) (13). Blood samples were taken from the tail vein 72 h after the STZ injection to measure the blood glucose levels by ACCU-Check glucose meter. Just animals with fasting blood glucose levels (after fasting for 12 h) over

300 mg/dL were considered diabetic and used for the further study (14).

Experimental design

All animals were randomly divided into eight groups with six animals in each group.

1. Normal control treated with normal saline (10 mL/Kg).

2. Normal rats treated with aqueous extract of *Berberis integerrima* root (250 mg/Kg body weight).

3. Normal rats treated with aqueous extract of *Berberis integerrima* root (500 mg/Kg body weight).

4. Normal rats treated with Glibenclamide (0.6 mg/Kg body weight)

5. Diabetic control treated with normal saline (10 mL/Kg).

6. Diabetic rats treated with aqueous extract of *Berberis integerrima* root (250 mg/Kg body weight).

7. Diabetic rats treated with aqueous extract of *Berberis integerrima* root (500 mg/Kg body weight).

8. Diabetic rats treated with Glibenclamide (0.6 mg/Kg body weight).

Animals were treated daily by gavage for 6 weeks and the experimental period for each rat was 6 weeks.

Blood collection

At the end of the study, animals were fasted overnight and anesthetized with chloroform (Pharmaceutical Partners of Japan). Blood samples were collected from the animal's hearts and the serum was separated by centrifugation (3000 rpm at 4°C for 15 min) and stored at -30°C for different biochemical analysis.

Estimation of body weight

The body weight in experimental animals was determined before the study and at 2, 4 and 6 weeks after it by a digital balance. These weights were determined at the same time during the morning.

Estimation of blood glucose

Throughout the 6-week treatment period, fasting (12 h) blood glucose was measured before the study, and at 2, 4 and 6 weeks after it on lateral

tail vein blood samples using an ACCU-Check glucose meter (Roche, Mannheim, Germany)

Estimation of some serum physiological parameters

All biochemical parameters in serum concentration including of triglycerides, cholesterol, high density lipoprotein HDL-cholesterol (HDL-C), low density lipoprotein LDL-cholesterol (LDL-C), total protein, creatinine, urea, alanine aminotransferase (ALT), aspartate aminotransferase (AST), alkaline phosphatase (ALP) and total bilirubin determined with the use of commercially available enzyme kits (Pars Azmoon, Tehran, Iran) and using an automatic analyzer (Architect c8000 Clinical Chemistry System, USA). LDL cholesterol (LDLC) was estimated by Frydvald method: $LDL\ cholesterol = total\ cholesterol - HDL\ cholesterol - (Triglyceride / 5)$.

Statistical analysis

All the reported data are expressed as mean \pm SEM. Statistical analysis was performed using one-way ANOVA followed by Tukey's multiple tests using 18th version of the computer software. The values were considered statistically significant when p-value was less than 0.05 compared to the respective control.

Results

Blood glucose

The hypoglycemic effect of AEBI and glibenclamide on the fasting blood sugar levels of normal and diabetic rats is shown in Table 1. The results clearly indicated that the AEBI (250 and 500 mg/Kg bw) or glibenclamide (0.6 mg/Kg bw) shows a significant hypoglycemic activity in normoglycemic rats by 22.48, 36.76, and 24.67% respectively in 2 weeks of treatment, however, at the end of 6th week, it was increased to achieve the normal blood sugar levels that are 86.40 ± 3.31 mg/dL, 94.00 ± 2.40 mg/dL and 90.80 ± 5.34 mg/dL respectively. The administration of STZ (65 mg/Kg, IP) led to about 3.5-fold elevation of fasting blood glucose levels. Diabetic rats treated with AEBI (250 and 500 mg/Kg bw) or glibenclamide (0.6 mg/Kg bw) showed a significant decrease in blood

Table 1. Effect of glibenclamide and AEBI on the blood glucose in normal and diabetic rats.

Group (n = 6)	Treatment	Dose (mg/Kg)	Blood glucose level (mg/dL)			
			Week 0	Week 2	Week 4	Week 6
1	N+C	10 mL/Kg	89.40 ± 5.98	89.20 ± 4.50	98.20 ± 4.52	93.80 ± 3.35
2	N+AEBI	250	89.00 ± 4.04	69.00 ± 4.04 ^b	77.40 ± 3.24 ^a	86.40 ± 3.31
3	N+AEBI	500	97.40 ± 2.48	61.60 ± 2.60 ^{b**}	82.20 ± 3.03	94.00 ± 2.40
4	N+G	0.6	89.20 ± 5.57	67.20 ± 3.02 ^{b*}	81.60 ± 4.44	90.80 ± 5.34
5	D+C	10 mL/Kg	89.40 ± 4.73	316.00 ± 2.16 ^{#**}	327.00 ± 3.92 ^{#**}	337.20 ± 7.53 ^{#**}
6	D+AEBI	250	97.20 ± 3.13	220.60 ± 3.09 ^{d**}	175.4 ± 5.50 ^{d**}	162.40 ± 2.61 ^{d**}
7	D+AEBI	500	89.80 ± 4.27	181.80 ± 2.26 ^{d**}	145.20 ± 2.74 ^{d**}	110.40 ± .92 ^d
8	D+G	0.6	87.20 ± 6.41	186.40 ± 2.37 ^{d**}	147.40 ± 3.60 ^{d**}	113.00 ± 3.43 ^{d**}

AEBI: Aqueous extract of *Berberis integerrima*; N: normal; C: control; G: glibenclamide; D: diabetic; Values are presented as mean ± SEM; n = 6 in each group. One way ANOVA followed by Tukey test. a p < 0.05 and bp < 0.01. Normal treated rats were compared with Normal control Rats. #p < 0.01 Diabetic control Rats were compared with Normal control Rats. cp < 0.05 and dp < 0.01 Diabetic treated rats were compared with Diabetic control rats on corresponding day; *p < 0.05 and **p < 0.01 compared to 0 value.

glucose level during the 6 weeks of treatment. The value of blood glucose levels of diabetic rats in the 6 weeks of treatment in comparison with diabetic control are 51.83, 67.25 and 66.48%, respectively.

Changes in body weight

As shown in Table 2, normal control animals were found to be stable in their body weight. Diabetic rats treated by 250 mg/Kg bw of AEBI showed a little reduction in body weight by percentage 3.84% during 6 weeks but diabetic rats treated by 500 mg/Kg bw of AEBI or glibenclamide (0.6 mg/Kg bw) showed a negligible increase in the body weight by 3.03 and 1.7% respectively. While normal rats treated by AEBI (250 and 500

mg/Kg bw) or glibenclamide (0.6 mg/Kg bw) showed a little reduction in the body weight by 16.02, 17.86 and 20.28% respectively, the body weight of diabetic control group was decreased to 33.3% during 6 weeks.

Serum lipid profiles

Table 3 shows the effect of aqueous root extracts and glibenclamide on serum lipid profiles of control and experimental groups. The hyperlipidemic parameters like serum triglyceride (TG), total cholesterol (TC), and LDL cholesterol (LDLC) were increased, but HDL cholesterol decreased in diabetic groups in comparison to the normal control. However, all these parameters except HDL cholesterol

Table 2. Effect of glibenclamide and AEBI on body weight in normal and diabetic rats.

Group (n = 6)	Treatment	Dose (mg/Kg)	Average body weight (g)			
			Week 0	Week 2	Week 4	Week 6
1	N+C	10 mL/Kg	193.20 ± 6.58	211.56 ± 4.74	233.34 ± 3.20 ^{**}	247.62 ± 1.87 ^{**}
2	N+AEBI	250	204.08 ± 6.03	216.28 ± 5.22	226.90 ± 4.78	236.78 ± 3.58 ^{**}
3	N+AEBI	500	200.94 ± 3.47	210.20 ± 2.91	219.76 ± 3.34	236.86 ± 3.18 ^{**}
4	N+G	0.6	197.86 ± 6.20	212.32 ± 4.41	223.64 ± 4.02 [*]	238.80 ± 2.41 ^{**}
5	D+C	10 mL/Kg	206.84 ± 3.35	155.28 ± 4.97 ^{#**}	144.90 ± 2.01 ^{#**}	137.98 ± 2.24 ^{#**}
6	D+AEBI	250	195.04 ± 4.67	183.44 ± 2.88 ^d	180.96 ± 3.74 ^d	187.56 ± 3.73 ^d
7	D+AEBI	500	194.36 ± 1.78	181.90 ± 1.62 ^d	183.52 ± 5.97 ^d	200.22 ± 2.78 ^d
8	D+G	0.6	199.94 ± 1.72	184.30 ± 4.08 ^d	193.46 ± 5.53 ^d	203.52 ± 5.61 ^d

AEBI: Aqueous extract of *Berberis integerrima*; N: normal; C: control; G: glibenclamide; D: diabetic; Values are presented as mean ± SEM. n = 6 in each group. One way ANOVA followed by Tukey test. a p < 0.05 and bp < 0.01 Normal treated rats were compared with Normal control rats. #p < 0.01 Diabetic control rats were compared with Normal control Rats. cp < 0.05 and dp < 0.01 Diabetic treated rats were compared with Diabetic control Rats on corresponding day; *p < 0.05 and **p < 0.01 compared to 0 value

Table 3. Effect of glibenclamide and AEBI on the lipid profiles (mg/dL) in normal and diabetic rats.

Group (n = 6)	Treatment	Dose (mg/Kg)	Serum lipid profiles (mg/dL)			
			TC	TG	HDL	LDL
1	N+C	10 mL/Kg	76.28 ± .98	66.42 ± 1.03	38.54 ± .48	24.25 ± 1.21
2	N+AEBI	250	74.40 ± .85	64.66 ± 1.79	39.95 ± .52	21.66 ± .93
3	N+AEBI	500	72.70 ± .54	60.96 ± .84	40.82 ± .61	19.68 ± .91
4	N+G	0.6	72.68 ± 1.02	64.26 ± .70	39.80 ± .74	19.58 ± 1.76
5	D+C	10 mL/Kg	125.94 ± 1.67 [#]	121.68 ± .90 [#]	14.46 ± .32 [#]	87.14 ± 1.63 [#]
6	D+AEBI	250	76.60 ± 1.55 ^d	85.26 ± 1.87 ^d	33.08 ± .76 ^d	26.46 ± 1.26 ^d
7	D+AEBI	500	69.48 ± 2.04 ^a ^d	73.84 ± 1.87 ^d	37.30 ± .65 ^d	17.41 ± 1.73 ^d
8	D+G	0.6	97.70 ± 2.49 ^b ^d	81.53 ± .94 ^d	35.20 ± .56 ^d	46.18 ± 2.67 ^d

AEBI: Aqueous Extract of *Berberis Integerrima*; N: normal; C: control; G: glibenclamide; D: diabetic; Values are presented as mean ± SEM. n = 6 in each group. One-way ANOVA followed by Tukey test. a p < 0.05 and bp < 0.01. Normal treated rats were compared with Normal control rats. #p < 0.01; Diabetic control rats were compared with Normal control rats. cp < 0.05 and dp < 0.01; Diabetic treated rats were compared with Diabetic control rats.

were decreased significantly in the diabetic groups treated by 250 and 500 mg/Kg of AEBI or glibenclamide (0.6 mg/Kg bw) (about 39.17, 44.82 and 22.42% for total cholesterol respectively), (about 29.93, 39.31, and 32.99% for triglyceride respectively) and (about 69.63, 80.02 and 47.00% for LDL cholesterol respectively) compared to diabetic rats. HDL cholesterol was increased significantly (about 128.76, 157.95 and 143.43% respectively) compared to diabetic rats. On the other hand, normal rats group treated by 250 and 500 mg/Kg bw of AEBI and glibenclamide (0.6 mg/Kg bw) for 6 weeks, caused a little significant reduction on all parameters (about 2.45, 4.69 and 4.71% for total cholesterol respectively), (about 2.62, 8.22 and 3.25% for Triglyceride respectively and about 10.68, 18.84 and 19.25% for LDL cholesterol respectively) compared to normal control; while, the HDL cholesterol was a negligible increased (about 3.65, 11.31 and 3.26% respectively) compared to the normal control.

Kidney parameters

The mean values of serum urea and creatinine concentrations of both control and experimental groups are presented in Table 4. STZ-induced diabetic rats showed a significant increase (p < 0.01) in blood urea and creatinine compared to normal control. Administration of 250 and 500 mg/Kg bw of AEBI or glibenclamide (0.6 mg/Kg) led to significant decrease in serum

creatinine and urea levels in STZ-induced diabetic rats (about 41.7, 52.94 and 45.29 for creatinine respectively and about 54.86, 65.56 and 62.26% for urea respectively) as compared with untreated STZ-induced diabetic rats. On the other hand normal rats treated with 250 and 500 mg/Kg bw of AEBI or glibenclamide (0.6 mg/Kg) for 6 weeks, caused a little reduction (about 3.03, 7.57 and 4.54% for creatinine respectively and about 7.04, 9.09 and 4.59% for urea respectively) compared to normal control.

Liver parameters

Table 5 shows the mean values of AST, ALT, ALP activities and serum total bilirubin and total protein levels of both control and experimental groups after 6 weeks. In STZ-induced diabetic rats, the activities of blood AST, ALT, ALP and the serum total bilirubin level were significantly increased (p < 0.01), but serum total protein levels were decreased (p < 0.01) compared to their normal levels. On the other hand, treatment of the STZ-induced diabetic rats by 250 and 500 mg/Kg of AEBI or glibenclamide (0.6 mg/Kg) caused a significant reduction in the activity of these parameters in blood by 42.52, 50.71 and 50.31%, respectively for AST and by 26.28, 50.36 and 42.64% respectively for ALT, and by 18.49, 43.82 and 37.52% respectively for ALP and by 25.89, 41 and 30.21% respectively for serum total bilirubin level but increased in blood total protein by 128.57, 267.34 and 134.69% compared to the mean values of untreated

Table 4. Effect of glibenclamide and AEBI on the liver parameters (mg/dL) in normal and diabetic rats.

Group (n = 6)	Treatment	Dose (mg/Kg)	liver parameters (mg/dL)				
			ALP	ALT	AST	T bilirubin	T protein
1	N+C	10 mL/Kg	93.82 ± 4.76	25.84 ± 1.60	21.00 ± 1.57	0.83 ± .04	2.06.08
2	N+AEBI	250	85.62 ± 4.15	23.28 ± 1.83	20.02 ± 0.99	0.74 ± .02	2.14 ± .22
3	N+AEBI	500	82.80 ± 1.59	21.10 ± 1.78	17.56 ± .99	0.76 ± .03	2.30 ± .23
4	N+G	0.6	84.06 ± 2.28	22.42 ± 2.08	19.10 ± .88	0.76 ± .03	2.17 ± .12
5	D+C	10 mL/Kg	180.78 ± 3.25 [#]	48.92 ± 1.96 [#]	44.68 ± 2.35 [#]	1.39 ± .06 [#]	0.49 ± .05 [#]
6	D+AEBI	250	147.34 ± 4.56 ^d	36.06 ± 1.89 ^d	25.68 ± 1.79 ^d	1.03 ± .07 ^d	1.15 ± .05 ^c
7	D+AEBI	500	101.56 ± 7.79 ^d	24.28 ± 1.74 ^d	22.02 ± 1.56 ^d	0.82 ± .05 ^d	1.80 ± .10 ^d
8	D+G	0.6	112.94 ± 5.48 ^d	28.06 ± 1.48 ^d	22.20 ± .65 ^d	0.97 ± .09 ^d	1.12 ± .08 ^c

AEBI: Aqueous Extract of Berberis Integerrima; N: normal; C: control; G: glibenclamide; D: diabetic; Values are presented as mean ± SEM; n = 6 in each group. One way ANOVA followed by Tukey test. a p < 0.05 and bp < 0.01; Normal treated rats were compared with Normal control rats. #p < 0.01; Diabetic control rats were compared with Normal control rats. cp < 0.05 and dp < 0.01; Diabetic treated rats were compared with diabetic control rats.

diabetic group. On the other hand, normal rats group treated by 250 and 500 mg/Kg bw of AEBI or glibenclamide (0.6 mg/Kg bw) for 6 weeks, caused a little reduction (about 4.66, 16.33 and 9.04 for AST respectively), (about 9.90, 18.34 and 13.23 for ALT respectively), (about 8.74, 11.74 and 10.40 for ALP respectively), (about 10.84, 8.43 and 8.43 for bilirubin respectively) and also a little increase in total protein (about 3.88, 11.65 and 5.33 respectively) compared to normal control.

Discussion

In the present study, the blood glucose level had significant increase in diabetic rats. This result is similar to the finding of Augusti *et al.* and Campos *et al.* in rats (15, 16), Kumar and Reddy in mice (17) and Jain and Vyas in rabbits (18). There are many studies in this regard that indicating the effective reduction in blood sugar levels in STZ-induced diabetic rats (19-21). In the present study, the AEBI (dose dependently) and glibenclamide (0.6 mg/Kg) showed a significant hypoglycemic activity in normoglycemic in 2 weeks of treatment, however, at the end of the 6th week, it was increased to achieve the normal blood sugar levels. Injection of STZ (65 mg/Kg, IP) led to about 3.5-fold elevation of fasting blood glucose levels. Diabetic rates treated by AEBI (dose dependently) or glibenclamide showed a significantly decrease in blood glucose level during 6 weeks of treatment in comparison

with diabetic control. The activity of this fraction could be due to the presence of Berberine, saponins, flavonoids, alkaloids and steroids components. Different mechanisms already exist to use the plant extracts for reducing the blood glucose. Based on the results, it is assumed that the root extract could be responsible for the stimulation insulin release and the observed restoration of metabolic activities. In addition, these results indicated that the extracts possess active phytochemical principles with either cytoprotective functions on the pancreatic β -cells or insulino-protective properties. The hypoglycemic activity in a number of other plants is applied through the stimulation of insulin release (22, 23). The hypoglycemic activity in some of other plants is similar to the well-known sulfonylurea drugs like glibenclamide. They reduce blood glucose in normoglycemic animals (24, 25). The glibenclamide effects on blood glucose levels by increasing the activity of pancreatic β -cells of the pancreas result in the secretion of a large amount of insulin (26, 27). The maximum reduction in serum glucose levels was seen in AEBI at dose of 500 mg/Kg (Table 1) hence, we could say that aqueous root extract of Berberis integerrima had a beneficial effect on carbohydrate.

In addition, results of the present study showed that diabetic rats exhibited a significant decrease in body weight after 6 weeks. These results are in agreement with those previously obtained (28-30). Body weight loss in diabetic rats may

Table 5. Effects of glibenclamide and AEBI on kidney parameters (mg/dL) in normal and diabetic rats.

Group (n = 6)	Treatment	Dose (mg/Kg)	Kidney parameters (mg/dL)	
			Serum creatinine	Serum urea
1	N+C	10 mL/Kg	0.66 ± .02	19.58 ± 1.24
2	N+AEBI	250	0.64 ± .02	18.20 ± 1.08
3	N+AEBI	500	0.61 ± .00	17.80 ± 1.37
4	N+G	0.6	0.63 ± .01	18.68±1.54
5	D+C	10 mL/Kg	1.70 ± .10 [#]	69.70±1.83 [#]
6	D+AEBI	250	1.00 ± .13 ^d	31.46±2.45 ^d
7	D+AEBI	500	0.80 ± .14 ^d	24.00±1.43 ^d
8	D+G	0.6	0.93 ± .06 ^d	26.30±1.36 ^d

AEBI: Aqueous Extract of *Berberis Integerrima*; N: normal; C: control; G: glibenclamide; D: diabetic; Values are presented as mean ± SEM. n = 6 in each group. One way ANOVA followed by Tukey test. a: p < 0.05 and b: p < 0.01 Normal treated rats were compared with Normal control rats. #p < 0.01; Diabetic control rats were compared with Normal control rats. c: p < 0.05 and d: p < 0.01; Diabetic treated rats were compared with Diabetic control rats.

be due to the fats and protein catabolism, even though the food intake is more in diabetic rats than control group. Due to the insulin deficiency protein, content is decreased in muscular tissue by proteolysis (31). Fat mobilization in skeletal muscle in STZ-induced diabetic rats (32) significantly induces the weight loss (33) as observed in the present study. However, diabetic rates treated by 250 and 500 mg/Kg of AEBI or glibenclamide (0.6 mg/Kg bw) showed a little reduction in body weight which may be explained by the increased insulin secretion or increased food consumption (34, 35).

Lipids play a major role in the disorder of diabetes mellitus. The most common lipid disorders in diabetes are hypertriglyceridemia and hypercholesterolemia (36). In our study, we have noticed significantly increased levels of serum total cholesterol, triglycerides and LDL-cholesterol but markedly decreased level of serum HDL-cholesterol in STZ-induced diabetic rats. These results are in agreement with those obtained (37-39). The main reason for the abnormal concentration of serum lipids in diabetic rats is an increase in the mobilization of free fatty acids from peripheral fat depots, since insulin inhibits the hormone-sensitive lipase (40). Excessed fatty acids in the serum of diabetic rats are converted into phospholipids and cholesterol in the liver. These two substances along with excessed triglycerides formed at the same time in the liver maybe discharged into the blood in the form of lipoproteins (41).

The present study showed that AEBI (250 and 500 mg/Kg) or glibenclamide (0.6 mg/Kg bw) had favorably modified serum lipid profile in rats with significant decreases (dose-dependently for AEBI) in total cholesterol, LDL-cholesterol, triglycerides and increased the serum HDL-C level in diabetic rats. These findings are consistent with earlier report that showed the administration of *Berberis vulgaris* Fruit Extract (3 g/Kg/d) for 3 months in type 2 diabetic patients decreased the Serum glucose, Triglycerides, total cholesterol and LDL-c and also increased the HDL-c (53).

The mechanism(s) of the hypolipidemic actions of AEBI are not known; however, they could be mediated by controlling tissue metabolism and improved insulin secretion and action since insulin lowers the lipid levels and normalizes lipids in STZ-induced diabetic rats (42).

In our study, we have noticed significantly increased levels of serum urea and creatinine in STZ-induced diabetic rats. Many studies have shown a significant increase in the rate of kidney cell damage (nephropathy) in diabetes disorders (43). Finally, this nephropathy reduces the physiological function and changes in the structure of kidney in diabetes (44). Hyperglycemia increases the generation of free radicals by glucose auto-oxidation and the increment of free radicals may lead to kidney cells damage (45). On the other hand, studies have shown that the increase in oxidative

stress the basement membrane of glomerular is thickening and the mesenchymal expanded (46). These changes lead to hypertrophy and decrease in physiological function of the kidney as well as increase in serum urea and creatinine levels (47). Our findings also show these changes. Previous changes in serum urea and creatinine concentrations strongly suggested impairment of kidney function in diabetes. The treatment with AEBI especially at dose of 500 mg/Kg and glibenclamide (0.6) return these parameters close to normal levels. The main effect of the AEBI and glibenclamide is presumably due to its ability to increase the insulin secretion since insulin lowers glucose levels and normalizes glucose in STZ-induced diabetic rats (22, 23).

The enzymes levels such as AST and ALT are largely used to assess the liver damage by streptozotocin. Necrosis or membrane damage releases the enzyme into circulation and hence it can be measured in the serum. High levels of AST indicates liver damage that occurs in many diseases such as viral hepatitis as well as cardiac infarction and muscle injury. AST catalyses the conversion of alanine to pyruvate and glutamate and is released in a similar manner. Therefore, ALT is more specific to the liver, and thus is a better parameter for detecting the liver injury. Elevated levels of serum enzymes indicate cellular leakage and functional integrity loss of cell membrane in liver (48). Serum ALP, bilirubin and total protein levels, on the other hand, are related to the function of hepatic cell. Increase in serum level of ALP is due to increased synthesis, in presence of increasing biliary pressure (49). Decline in serum total protein may be due to the inhibited oxidative phosphorylation process which leads to a decrease in protein synthesis, an increase in the catabolic process and reduction of protein absorption (50). STZ-induced caused a significant ($p < 0.001$) elevation in enzyme levels such as AST, ALT, ALP, level total bilirubin and decrease in total protein when compared to control. There was a significant ($p < 0.001$) restoration of these enzyme levels on the administration of AEBI in a dose-dependent manner and also by glibenclamide at dose of 0.6 mg/Kg. The reversal of increased serum enzymes in STZ-induced liver damage by the

extract may be due to the prevention of the leakage of intracellular enzymes by its membrane stabilizing activity. This is in agreement with the commonly accepted view that serum levels of transaminases return to normal with the healing of hepatic parenchyma and the regeneration of hepatocytes (51). Effective control of ALP, bilirubin and total protein levels point toward an early improvement in the secretory mechanism of the hepatic cells. The efficacy of any hepatoprotective drug is dependent on its capacity of either reducing the harmful effect or restoring the normal hepatic physiology that has been distributed by a hepatotoxin. Both glibenclamide and the aqueous extract of *Berberis integerrima* root decreased the STZ-induced elevated enzyme levels in tested groups, indicating the protection of structural integrity of hepatocytic cell membrane or regeneration of damaged liver cells.

Conclusion

The aqueous extract of *Berberis integerrima* root exhibited significant antihyperglycemic, antihyperlipidemic and antioxidant activities in STZ-induced diabetes. These effects of root extract of *Berberis integerrima* especially at dose of 500 mg/Kg in all parameters except for the blood glucose, (similar) are more than glibenclamide in diabetic rats. The results support the use of this extract in traditional medicine, although the total extract does not confer any additional benefits or disadvantages compared with the well-known sulfonylurea drugs like glibenclamide.

Acknowledgment

Authors would like to express thanks to Ms. Farnad of Biochemistry Laboratory Expert, Biology Department, Urmia University, Iran, for providing laboratory facilities.

References

- (1) Okoli CO, Ibiam AF, Ezike AC, Akah PA and Okoye TC. Evaluation of antidiabetic potentials of *Phyllanthus niruri* in alloxan diabetic rats. *Afr. J. Biotechnol.* (2010) 9: 248-59.

- (2) Wild S, Roglic G, Green A, Sicree R and King H. Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes Care*. (2004) 27: 1047-53.
- (3) Esteghamati A, Gouya MM, Abbasi M, Delavari A, Alikhani S and Alaedini F. Prevalence of diabetes and impaired fasting glucose in the adult population of Iran: National Survey of Risk Factors for Non Communicable Diseases of Iran. *Diabetes-care*. (2008) 31: 96-98.
- (4) Suji G and Sivakami S. Approaches of diabetes mellitus: an overview. *Cell Mol. Biol. (Noisy-le-grand)* (2003) 49: 635-9.
- (5) Tripathi BK and Srivastava AK. Diabetes mellitus : complications and therapeutics. *Med. Sci-Monit* (2006) 12: 130-47.
- (6) Shapiro K and Gong WC. Natural products used for diabetes. *J. Am. Pharm. Assoc. (Wash)* (2002) 42: 217-26.
- (7) Arayne M S, Sultana N and SherBahadur S. The berberis story: *Berberis vulgaris* in therapeutics. *Pak. J. Sci.* (2007) 20: 83-92.
- (8) Wong F and YiZhou. Inhibitory effects of berberine on ion channels of rat hepatocytes. *World. J. Gastroenterol.* (2004) 10: 2842-2845.
- (9) Issat T and Jakobisiak M. Berberine a natural cholesterol reducing product, exerts antitumor cytostatic/ cytotoxic effects independently from the mevalonate pathway. *Oncology reports. Golab. J.* (2006) 16: 1273-1276.
- (10) Vrzal R, Zdarilova A and Ulrichova J. Activation of the aryl hydrocarbon receptor by berberin in H4IIE and HepG2 cells. *Science direct. doi:10.1016/J.bcp.* (2005) 925-936.
- (11) Nawel M, Mohamed E, Amine D, Hocine A, Boufeldja T. Hypoglycaemic effect of *Berberis vulgaris* L. in normal and streptozotocin-induced diabetic rats. *Asian Pac. J. Trop. Biome.* (2011) 468-471.
- (12) OECD 2002. Acute oral toxicity. Acute oral toxic class method guideline 423 adopted 23.03.1996. In: Eleventh Addendum to the, OECD, guidelines for the testing of chemicals organization for economical cooperation and development, Paris, June (2000).
- (13) Sancheti S, Sancheti S, Bafna M, Seo SY. Antihyperglycemic, antihyperlipidemic, and antioxidant effects of *Chaenomeles sinensis* fruit extract in streptozotocin induced diabetic rats. *Eur. Food Res. Technol.* (2010) 231: 415.
- (14) Hosseinzadeh H, Ramzani M and Danaei AR. Antihyperglycemic effect and acute toxicity of *Securiger a securidaca* L. seed Extracts in mice. *Phytotherapy Res.* (2002) 16: 745-7.
- (15) Augusti K and Sheela C. Antiperoxide effect of S-allyl cysteine sulfoxide, an insulin secretagogue, in diabetic rats. *Experientia* (1996) 52: 115-120.
- (16) Campos K, Diniz Y, Cataneo A, Faine L, Alves J and Novelli E. Hypoglycaemic and antioxidant effects of onion, *Allium cepa*: dietary onion addition, antioxidant activity and hypoglycaemic effects on diabetic rats. *Int. J. Food Sci. Nutr.* (2003) 54: 241-246.
- (17) Kumar G and Reddy K. Reduced nociceptive responses in mice with alloxan induced hyperglycemia after garlic (*Allium sativum*) treatment. *Indian J. Exp. Biol.* (1999) 37: 662-666.
- (18) Jain R and Vyas C. Garlic in alloxan- induced diabetic rabbits. *Am. J. Clin. Nutr.* (1975) 28: 684-685.
- (19) Ravi K, Ramachandra B and Subramanian S. Protective effect of *Eugenia jambolana* seed kernel on tissue antioxidants in streptozotocin induced diabetic rats. *Biol. Pharm. Bull.* (2004) 27: 1212-1217.
- (20) Rajasekaran S, Sivagnanam K and Subramanian S. Antioxidant effect of *Aloe vera* gel extract in streptozotocin-induced diabetes in rats. *Pharmacol. Reports.* (2005) 57: 90-96.
- (21) Sekar D, Sivagnanam K and Subramanian S. Antidiabetic activity of *Momordica charantia* seeds on streptozotocin induced diabetic rats. *Pharmazie* (2005) 60: 383-7.
- (22) Kumar S, Kumar V and Prakash OM. Antidiabetic and anti-lipemic effects of *Cassia siamea* leaves extract in streptozotocin induced diabetic rats. *Asian. Pac. J. Trop. Med.* (2010) 3: 871-873.
- (23) Shirwaikar A, Rajendran K and Punitha ISR. Antidiabetic activity of alcoholic stem extract of *Coscinium fenestratum* in streptozotocin-induced type 2 diabetic rats. *J. Ethnopharmacol.* (2005) 97: 369-374.
- (24) Jain S, Bhatia G, Barik R, Kumar P, Jain A and Dixit VK. Antidiabetic activity of *Paspalum scrobiculatum* Linn. in alloxan induced diabetic rats. *J. Ethnopharmacol.* (2010) 127: 325-328.
- (25) Nagarajana NS, Muruges N, Thirupathy Kumaresan P, Radha N and Murali A. Antidiabetic and antihyperlipemic effects of *Cleomea felina*. *Fitoterapia* (2005) 76: 310-315.
- (26) Shalev A. Hope for insulin mimetic oral antidiabetic drugs. *Eur. J. Endocrinol.* (1999) 8: 561-562.
- (27) Ozturk Y, Altan VM and Yildizoglu A. Effect of experimental diabetes and insulin on smooth muscle functions. *Pharmacol. Rev.* (1996) 10: 69-112.
- (28) Ananthi J, Prakasam A and Pugalendi K. Antihyperglycemic activity of *Eclipta alba* leaf on alloxan-induced diabetic rats. *Yale. J. Biol. Med.* (2003) 76: 97- 102.
- (29) Howarth F, Jacobson M, Naseer O and Adeghate E. Short-term effects of streptozotocin-induced diabetes on the electrocardiogram, physical activity and body temperature in rats. *Exp. Physiol.* (2004) 90: 237-245.
- (30) AL-Rawi M. Effect of *Trifolium* sp. flowers extracts on the status of liver histology of streptozotocin-induced diabetic rats. *Saudi J Bio Sci* (2007) 14:21-28.
- (31) Vats V, Yadav S and Grover J. Ethanolic extract of *Ocimum sanctum* leaves partially attenuates streptozotocin-induced alterations in glycogen content and carbohydrate metabolism in rats. *J. Ethnopharmacol.* (2004) 90: 155-160.
- (32) Stearns S, Tepperman H and Tepperman J. Studies on the utilization and mobilization of lipid in skeletal muscles from streptozotocin-diabetic and control rats.

- J. Lipid. Res.* (1979) 20: 654-662.
- (33) Besse S, Assayag P, Delcayre C, Carré F, Cheav SL, Lecarpentier Y and Swynghedauw B. Normal and hypertrophied senescent rat heart: mechanical and molecular characteristics. *Am. J. Physiol.* (1993) 265: H183-H190.
- (34) Farouque H, Meredith L. Effects of inhibition of ATP sensitive potassium channels on metabolic vasodilation in the human forearm. *Clin. Sci.* (2003) 104: 39-46.
- (35) Fernstrom M, Fernstrom J. Large changes in serum free tryptophan levels do not alter tryptophan levels. Studies in streptozotocin, diabetic rats. *Life. Sci.* (1993) 52: 907- 916.
- (36) Al-Shamaony L, Al-khazraji S and Twaij H. Hypoglycaemic effect of Artemisia herba alba. II. Effect of a valuable extract on some blood parameters in diabetic animals. *J. Ethnopharmacol* (1994) 43: 167-171.
- (37) Bolkent S, Yanardag R, Karabulut-Bulen O and Ozsoy-Sacan O. The morphological and biochemical effects of glibornuride on rat liver in experimental diabetes. *Hum. Exp. Toxicol.* (2004) 23: 257-264.
- (38) Singh N, Kamath V, Rajini P. Protective effect of potato peel powder in ameliorating oxidative stress in streptozotocin diabetic rats. *Plant. Foods. Hum. Nutr.* (2005) 60: 49-54.
- (39) Rajasekaran S, Ravi K, Savgananam K, Subramanian S. Beneficial effects of aloe vera leaf gel extract on lipid profile status in rats with streptozotocin diabetes. *Clin. Exp. Pharmacol. Physiol.* (2006) 33: 232-237.
- (40) Pushparaj P, Tan C, Tan B. Effects of Averrhoa bilimbi leaf extract on blood glucose and lipids in streptozotocin diabetic rats. *J. Ethnopharmacol.* (2000) 72: 69- 76.
- (41) Bopanna K, Kannan J, Sushma G, Balaram R and Rathod S. Antidiabetic and antihyperlipaemic effects of neem seed kernel powder on alloxan diabetic rabbits. *Indian J. Pharmacol.* (1997) 29: 162-167.
- (42) Pepato MT, Mori DM, Baviera AM, Harami JB, Vendramini RC and Brunetti IL. Fruit of the jambolan tree (*Eugenia jambolana* Lam.) and experimental diabetes. *J. Ethnopharmacol.* (2005) 96: 43-8.
- (43) Ahmed MH and Osman MM. Improving laboratory diagnosis of diabetic nephropathy with the use of glomerular filtration rate. *Diabetes Technol. Ther.* (2006) 8: 688-90.
- (44) Rosolowsky ET, Niewczas MA, Ficociello LH, Perkins BA, Warram JH and Krolewski AS. Between hyperfiltration and impairment: demystifying early renal functional changes in diabetic nephropathy. *Diabetes Res. Clin. Pract.* (2008) 13: Suppl 1: S46-53.
- (45) Sharma S, Kulkarni SK, Chopra K. Curcumin, the active principle of turmeric (*Curcuma longa*), ameliorates diabetic nephropathy in rats. *Clin. Exp. Pharmacol. Physiol.* (2006) 33: 940-5.
- (46) Jäckle-Meyer I, Szukics B, Neubauer K, Metzke V, Petzoldt R, Stolte H. Extracellular matrix proteins as early markers in diabetic nephropathy. *Eur. J. Clin. Chem. Clin. Biochem.* (1995) 33: 211-9.
- (47) Marice AD, Abd-Allah GM and El-Yamany MF. Renal oxidative stress and nitric oxide production in streptozotocin-induced diabetic nephropathy in rats: the possible modulatory effects of garlic (*Allium sativum* L.). *Biotechnol. Appl. Biochem.* (2009) 52: 227-32.
- (48) Drotman R and Lawhan G. Serum enzymes are indications of chemical induced liver damage. *Drug Chem. Toxicol.* (1978) 1: 163-171.
- (49) Muriel P and Garcipiana T. Silymarin protects against paracetamol-induced lipid peroxidation and liver damage. *J. Appl. Toxicol.* (1992) 12: 439-442.
- (50) Jefferson L, Liao W, Peavy D, Miller, Appel M and Taylor J. Diabetes-induced alterations in liver protein synthesis: Changes in the relative abundance of mRNA for albumin and other plasmaproteins. *J. Biol. Chem.* (1983) 258: 1369-1375.
- (51) Thabrew M and Joice P A. comparative study of the efficacy of Pavetta indica and Osbeckia octanda in the treatment of liver dysfunction. *Planta. Med.* (1987) 53: 239-241.
- (52) Minaiyan M, Ghannadi AR, Mahzouni P and Jaffari-Shirazi E. Comparative study of Berberis vulgaris fruit extract and berberine chloride effects on acetic acid-induced colitis in rats. *Iranian J. Pharm. Res.* (2011) 10: 97-104.
- (53) Shidfar F, Seyyed Ebrahimi S, Hosseini S, Heydari I, Shidfar S and Hajhassani G. The Effects of *Berberis vulgaris* Fruit Extract on Serum Lipoproteins, apoB, apoA-I, Homocysteine, Glycemic Control and Total Antioxidant Capacity in Type 2 Diabetic Patients. *Iranian J. Pharm. Res.* (2012) 11 643-652.