

Synthesis and Antimycobacterial Activity of Symmetric Thiocarbohydrazone Derivatives against *Mycobacterium bovis* BCG

Kamaleddin Haj Mohammad Ebrahim Tehrani^a, Farzad Kobarfard^{a,b}, Parisa Azerang^c, Maryam Mehravar^c, Zohreh Soleimani^a, Ghazaleh Ghavami^c and Soroush Sardari^{c*}

^aDepartment of Medicinal Chemistry, School of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran. ^bPhytochemistry Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran. ^cDrug Design and Bioinformatics Unit, Medical Biotechnology Department, Biotechnology Research Center, Tehran, Pasteur Institute, 13164, Iran.

Abstract

In this work, we reported the synthesis and evaluation of antimycobacterial and antifungal activity of a series of thiocarbohydrazone derivatives which are thiacetazone congeners. The target compounds were synthesized in superior yields by reacting thiocarbohydrazone with different aromatic aldehydes and methyl ketones. Compounds 8, 19 and 25 were found to be the most potent derivatives, exhibiting acceptable activity against *Mycobacterium bovis* BCG compared to thiacetazone and ethambutol as reference substances. Compounds 8, 15 and 25 exhibited the highest activity against *Candida albicans*. The most active compounds had a completely different aromatic ring system with various electronic, steric and lipophilic natures. This is understandable in light of the fact that carbohydrazone derivatives must undergo a metabolic activation step before exerting their anti-TB activity and different SAR rules govern each one of these two processes.

Keywords: Thiocarbohydrazone; Thiacetazone; *Mycobacterium bovis*; Antifungal; *In silico*.

Introduction

Tuberculosis (TB) is one of the most prevalent infectious diseases and based on the World Health Organization (WHO) report, 196 countries reported 2.6 million new positive TB cases in 2008 and among them, 1.78 million individuals died from TB (1). Although there are many on-going activities by pharmaceutical companies and researchers to fight the disease, there is still no great progress in this regard and emergence of resistance to the current antituberculosis drugs is widespread.

A serious concern in antitubercular therapy is the emergence of multi-drug resistant (MDR) strains, and more recently, extensively drug-resistant (XDR) strains of *Mycobacterium tuberculosis* (2). Based on a recent WHO report, 10% of MDR cases were XDR, across all geographical regions surveyed, and thus posing the threat of an untreatable global epidemic (3, 4). Therefore, there is a need for rapid and continued progress in development of new antitubercular agents and the discovery of new cellular drug targets.

Thiacetazone (TAC) is an economical, antitubercular, bacteriostatic drug that has been widely used in combination with other antimycobacterial agents like isoniazid (5).

* Corresponding author:

E-mail: ssardari@hotmail.com




Figure 1. Structures of SRI-224, SRI-286 and KBF611.




Figure 2. (a) Thiocetazone, (b) General structure of the new derivatives.

Molecular analogues of TAC, SRI-224 and SRI-286, have been synthesized and tested against *Mycobacterium avium* and found to be more effective than TAC *in-vitro* and *in-vivo* (mice) (6). In addition, our previous work on the halogenated analogues of thiocetazone led to the discovery of 4-acetamido-2-fluorobenzaldehyde thiosemicarbazone (KBF611), which was both non-toxic and highly active against *M. tuberculosis* (7). It has been shown that TAC is a prodrug that is activated by the mycobacterial monoxygenase EthA, which is also the activator of two other anti-tuberculosis drugs, ethionamide (ETH) and isoxyl (ISO) (8-10). However, the mechanism of action of TAC remains an enigma. There are indications that TAC affects mycolic acid synthesis in *M. bovis* BCG (8).

Structure activity relationship (SAR) studies on TAC shows that the thiosemicarbazone moiety is essential for antituberculosis activity. Our previous study showed the repeated presence of such molecular fragment category in compounds library with potential antimycobacterial activity (11). Therefore, we decided to synthesize a group of compounds which could be considered as a dimerized version of TAC. In these analogs, a thiocarbohydrazone moiety is linking two aromatic systems with different substitutions, various lipophilicity, electronic and steric characteristics.

Experimental

General

Melting points were obtained by an Electrothermal 9100 apparatus and are uncorrected. Infrared spectra were determined with a Perkin-Elmer 843 spectrometer. Proton nuclear magnetic resonance (^1H NMR) spectra and carbon nuclear magnetic resonance (^{13}C NMR) spectra were determined on a Bruker Avance DRX 500 MHz spectrometer and chemical shifts are reported as δ (ppm) in $\text{DMSO-}d_6$ solution (0.05% v/v TMS). ESI-MS spectra were obtained using Agilent 6410 Triple Quad. LC/MS. All the compounds were analyzed for C, H, N and S on a Costech model 4010 and agreed with the proposed structures within $\pm 0.4\%$ of the theoretical values.

General procedure for preparation of thiocarbohydrazones

To a hot solution of thiocarbohydrazide (0.21 g, 0.002 mol) in water (6 mL) containing acetic acid (0.4 mL), selected aldehydes or methyl ketones (0.004 mol) dissolved in ethanol (15 mL) were added dropwise and the mixture was heated under reflux. After the completion of the reaction, which was determined by thin layer chromatography, the reaction was cooled to room temperature and the precipitate thus formed was

filtered and rinsed with cold ethanol. In the cases that purification was needed, the crudes were recrystallized from appropriate solvents.

Bis(benzaldehyde) thiocarbohydrazone (1)

White powder (0.50 g, 89%): mp 195-198°C; IR (KBr): 3303, 3175, 1606, 1542, 1525, 1254, 1130, 774, 703 cm^{-1} ; ^1H NMR (DMSO- d_6 /500 MHz): δ 7.42-7.48 (m, 6H, aromatic H-3,4,5,3',4',5'), 7.76 (br s, 2H, aromatic H-1,5), 7.87 (br s, 2H, aromatic H-1',5'), 8.16 (br s, 1H, azomethine H), 8.62 (br s, 1H, azomethine H), 11.58 (br s, 1H, N-H), 11.90 (br s, 1H, N-H); ^{13}C NMR (DMSO- d_6 /125 MHz): 127.29, 128.69, 130.01, 134.12, 143.46, 148.76, 174.85; MS (ESI): 283 (M + H⁺), 305 (M + Na⁺); Anal. Calcd for C₁₅H₁₄N₄S (282.36): C, 63.80; H, 5.00; N, 19.84; S, 11.36. Found: C, 63.21; H, 4.99; N, 19.76; S, 11.32.

Bis(4-acetamidobenzaldehyde) thiocarbohydrazone (2)

Yellow powder (0.70 g, 88%): mp 188-190°C; IR (KBr): 3278, 3184, 1683, 1612, 1537, 1522, 1183, 849 cm^{-1} ; ^1H NMR (DMSO- d_6 /500 MHz): δ 2.06 (s, 6H, methyl H), 7.67 (d, $J = 6.65$, 4H, aromatic H-3,5,3',5'), 7.67 (br s, 2H, aromatic H-2,6), 7.79 (br s, 2H, aromatic H-2',6'), 8.07 (br s, 1H, azomethine H), 8.53 (br s, 1H, azomethine H), 10.13 (s, 2H, acetamide N-H), 11.42 (br s, 1H, thiocarbamide N-H), 11.77 (br s, 1H, thiocarbamide N-H); ^{13}C NMR (DMSO- d_6 /125 MHz): 24.04, 118.84, 128.00, 140.94, 142.95, 148.39, 168.53, 174.30; MS (ESI): 419 (M + Na⁺); Anal. Calcd for C₁₉H₂₀N₆O₂S (396.47): C, 57.56; H, 5.08; N, 21.20; S, 8.09. Found: C, 57.62; H, 5.11; N, 21.17; S, 8.12.

Bis(2-fluorobenzaldehyde) thiocarbohydrazone (3)

White powder (0.5 g, 79%): mp 196-199°C; IR (KBr): 3288, 3122, 1612, 1537, 1518, 1252, 771 cm^{-1} ; ^1H NMR (DMSO- d_6 /500 MHz): δ 7.32 (m, 4H, aromatic H-4,5,4',5'), 7.50 (m, 2H, aromatic H-3,3'), 7.98 (br s, 1H, aromatic H-6), 8.35 (br s, 1H, aromatic H-6'), 8.39 (br s, 1H, azomethine H), 8.88 (br s, 1H, azomethine H), 11.81 (br s, 1H, N-H), 12.08 (br s, 1H, N-H); ^{13}C NMR (DMSO- d_6 /125 MHz): 115.78, 115.94, 121.65, 124.73, 126.71, 131.94, 136.08, 141.41,

159.86, 161.85, 175.17; MS (ESI): 319 (M + H⁺), 341 (M + Na⁺); Anal. Calcd for C₁₅H₁₂F₂N₄S (318.34): C, 56.59; H, 3.80; N, 17.60; S, 10.07. Found: C, 56.71; H, 3.81; N, 17.58; S, 10.02.

Bis(3-fluorobenzaldehyde) thiocarbohydrazone (4)

White powder (0.51 g, 80%): mp 206-207°C; IR (KBr): 3293, 3129, 1580, 1547, 1517, 1259, 1138, 876, 763, 689 cm^{-1} ; ^1H NMR (DMSO- d_6 /500 MHz): δ 7.27 (dt, $J = 8.5$, $J = 2.2$, 2H, aromatic H-5,5'), 7.52 (m, 3H, aromatic H-6,4,4'), 7.59 (d, $J = 7.55$, 2H, aromatic H-2,2'), 7.91 (br s, 1H, aromatic H-6'), 8.15 (br s, 1H, azomethine H), 8.62 (br s, 1H, azomethine H), 11.73 (br s, 1H, N-H), 12.05 (br s, 1H, N-H); ^{13}C NMR (DMSO- d_6 /125 MHz): 112.81, 112.99, 116.71, 116.88, 123.62, 124.64, 130.78, 136.72, 142.16, 147.57, 161.46, 163.40, 175.14; MS (ESI): 319 (M + H⁺), 341 (M + Na⁺); Anal. Calcd for C₁₅H₁₂F₂N₄S (318.34): C, 56.59; H, 3.80; N, 17.60; S, 10.07. Found: C, 56.39; H, 3.81; N, 17.53; S, 10.11.

Bis(4-fluorobenzaldehyde) thiocarbohydrazone (5)

White powder (0.52 g, 82%): mp 224-227°C; IR (KBr): 3264, 3150, 1604, 1550, 1235, 1156, 843 cm^{-1} ; ^1H NMR (DMSO- d_6 /500 MHz): δ 7.31 (t, $J = 8.43$, 4H, aromatic H-3,5,3',5'), 7.81 (br s, 2H, aromatic H-2,6), 7.94 (br s, 2H, aromatic H-2',6'), 8.14 (br s, 1H, azomethine H), 8.59 (br s, 1H, azomethine H), 11.61 (br s, 1H, N-H), 11.90 (br s, 1H, N-H); ^{13}C NMR (DMSO- d_6 /125 MHz): 115.68, 115.86, 129.49, 130.73, 124.16, 147.57, 162.11, 164.08, 174.84; MS (ESI): 319 (M + H⁺), 341 (M + Na⁺); Anal. Calcd for C₁₅H₁₂F₂N₄S (318.34): C, 56.59; H, 3.80; N, 17.60; S, 10.07. Found: C, 56.82; H, 3.79; N, 17.56; S, 10.10.

Bis(2-chlorobenzaldehyde) thiocarbohydrazone (6)

White powder (0.67 g, 95%): mp 207°C (dec.); IR (KBr): 3301, 3192, 1603, 1539, 1478, 1247, 775 cm^{-1} ; ^1H NMR (DMSO- d_6 /500 MHz): δ 7.46 (m, 4H, aromatic H-4,5,4',5'), 7.52 (m, 2H, aromatic H-3,3'), 8.06 (br s, 1H, aromatic H-6), 8.39 (br s, 1H, aromatic H-6'), 8.61 (br s, 1H, azomethine H), 8.99 (br s, 1H, azomethine

H), 11.97 (br s, 1H, N-H), 12.15 (br s, 1H, N-H); ^{13}C NMR (DMSO- d_6 /125 MHz): 127.48, 127.80, 129.90, 131.52, 133.37, 139.73, 144.63, 175.37; MS (ESI): 373 (100%), 375 (50), 377 (15%) (M + Na⁺); Anal. Calcd for C₁₅H₁₂Cl₂N₄S (351.25): C, 51.29; H, 3.44; N, 15.95; S, 9.13. Found: C, 51.11; H, 3.45; N, 15.89; S, 9.08.

Bis (3-chlorobenzaldehyde) thiocarbohydrazone (7)

White powder (0.67 g, 95%): mp 216-219°C; IR (KBr): 3300, 3122, 1549, 1524, 1257, 1155, 893, 798, 695 cm⁻¹; ^1H NMR (DMSO- d_6 /500 MHz): δ 7.50 (s, 4H, aromatic H-4,5,4',5'), 7.71 (s, 2H, aromatic H-6,6'), 7.79 (br s, 1H, aromatic H-1), 8.11 (br s, 2H, aromatic H-1'), azomethine H), 8.60 (br s, 1H, azomethine H), 11.78 (br s, 1H, N-H), 12.07 (br s, 1H, N-H); ^{13}C NMR (DMSO- d_6 /125 MHz): 126.17, 126.93, 129.69, 130.61, 133.70, 136.10, 141.97, 147.20, 175.16; MS (ESI): 373 (100%), 375 (71%), 377 (17%) (M + Na⁺); Anal. Calcd for C₁₅H₁₂Cl₂N₄S (351.25): C, 51.29; H, 3.44; N, 15.95; S, 9.13. Found: C, 51.29; H, 3.42; N, 16.01; S, 9.09.

Bis (4-chlorobenzaldehyde) thiocarbohydrazone (8)

White powder (0.53 g, 75%): mp 205°C (dec.); IR (KBr): 3198, 3127, 1587, 1542, 1508, 1244, 825 cm⁻¹; ^1H NMR (DMSO- d_6 /500 MHz): δ 7.53 (d, $J = 8.32$, 4H, aromatic H-3,5,3',5'), 7.77 (br s, 2H, aromatic H-2,6), 7.91 (br s, 2H, aromatic H-2',6'), 8.14 (br s, 1H, azomethine H), 8.60 (br s, 1H, azomethine H), 11.67 (br s, 1H, N-H), 11.99 (br s, 1H, N-H); ^{13}C NMR (DMSO- d_6 /125 MHz): 128.87, 129.29, 132.90, 133.25, 134.56, 142.15, 147.48, 174.95; MS (ESI): 351 (100%), 353 (50%), 355 (11%) (M + H⁺), 373 (100%), 375 (77%), 377 (20%) (M + Na⁺); Anal. Calcd for C₁₅H₁₂Cl₂N₄S (351.25): C, 51.29; H, 3.44; N, 15.95; S, 9.13. Found: C, 51.11; H, 3.44; N, 16.00; S, 9.10.

Bis (2-bromobenzaldehyde) thiocarbohydrazone (9)

White powder (0.73 g, 83%): mp 208°C (dec.); IR (KBr): 3275, 3101, 1534, 1513, 1245, 1148, 685 cm⁻¹; ^1H NMR (DMSO- d_6 /500 MHz): δ 7.38 (dt, $J = 7.63$, $J = 1.70$, 2H, aromatic H-4,4'), 7.50 (t, $J = 7.63$, 2H, aromatic H-5,5'),

7.70 (dd, $J = 8.03$, $J = 0.66$, aromatic H-3,3'), 8.04 (br s, 1H, aromatic H-6), 8.37 (br s, 1H, aromatic H-6'), 8.58 (br s, 1H, azomethine H), 8.93 (br s, 1H, azomethine H), 12.03 (br s, 1H, N-H), 12.18 (br s, 1H, N-H); ^{13}C NMR (DMSO- d_6 /125 MHz): 123.81, 127.97, 131.76, 132.69, 133.15, 142.18, 146.87, 175.41; MS (ESI): 439 (60%), 441 (100%), 443 (50%) (M + H⁺), 461 (60%), 463 (100%), 465 (50%) (M + Na⁺); Anal. Calcd for C₁₅H₁₂Br₂N₄S (440.16): C, 40.93; H, 2.75; N, 12.73; S, 7.28. Found: C, 41.01; H, 2.78; N, 12.69; S, 7.25.

Bis (3-bromobenzaldehyde) thiocarbohydrazone (10)

White powder (0.76 g, 86%): mp 225°C (dec.); IR (KBr): 3285, 3142, 1528, 1515, 1245, 761 cm⁻¹; ^1H NMR (DMSO- d_6 /500 MHz): δ 7.42 (t, $J = 7.79$, 2H, aromatic H-5,5'), 7.63 (d, $J = 7.79$, 2H, aromatic H-4,4'), 7.75 (br s, 2H, aromatic H-6,6'), 7.92 (br s, 1H, aromatic H-1), 8.12 (br s, 1H, aromatic H-1'), 8.22 (br s, 1H, azomethine H), 8.58 (br s, 1H, azomethine H), 11.79 (br s, 1H, N-H), 12.05 (br s, 1H, N-H); ^{13}C NMR (DMSO- d_6 /125 MHz): 122.14, 126.79, 128.98, 130.77, 132.48, 136.46, 141.84, 146.90, 175.13; MS (ESI): 439 (55%), 441 (100%), 443 (56%) (M + H⁺); Anal. Calcd for C₁₅H₁₂Br₂N₄S (440.16): C, 40.93; H, 2.75; N, 12.73; S, 7.28. Found: C, 40.85; H, 2.75; N, 12.71; S, 7.26.

Bis (4-bromobenzaldehyde) thiocarbohydrazone (11)

White powder (0.82 g, 93%): mp 214°C (dec.); IR (KBr): 3190, 1581, 1538, 1245, 1114, 821 cm⁻¹; ^1H NMR (DMSO- d_6 /500 MHz): δ 7.67 (d, $J = 7.50$, 4H, aromatic H-3,5,3',5'), 7.67 (br s, 2H, aromatic H-2,6), 7.83 (br s, 2H, aromatic H-2',6'), 8.12 (br s, 1H, azomethine H), 8.58 (br s, 1H, azomethine H), 11.68 (br s, 1H, N-H), 11.99 (br s, 1H, N-H); ^{13}C NMR (DMSO- d_6 /125 MHz): 123.38, 128.98, 129.51, 131.81, 133.22, 133.58, 142.28, 147.61, 174.94; MS (ESI): 439 (40%), 441 (100%), 443 (40%) (M + H⁺), 461 (45%), 463 (100%), 465 (67%) (M + Na⁺); Anal. Calcd for C₁₅H₁₂Br₂N₄S (440.16): C, 40.93; H, 2.75; N, 12.73; S, 7.28. Found: C, 41.10; H, 2.77; N, 12.72; S, 7.26.

Bis(2-nitrobenzaldehyde) thiocarbohydrazone (12)

Yellow powder (0.66 g, 89%): mp 215°C (dec.); IR (KBr): 3263, 3105, 1506, 1339, 1237, 1112, 747 cm⁻¹; ¹H NMR (DMSO-*d*₆/500 MHz): δ 7.70 (dt, *J* = 7.64, *J* = 1.30, 2H, aromatic H-4,4'), 7.84 (t, *J* = 7.64, 2H, aromatic H-5,5'), 8.08 (dd, *J* = 7.64, *J* = 0.90, 2H, aromatic H-3,3'), 8.18 (br s, 1H, aromatic H-6), 8.43 (br s, 1H, aromatic H-6'), 8.63 (br s, 1H, azomethine H), 8.98 (br s, 1H, azomethine H), 12.10 (br, s, N-H), 12.30 (br, s, N-H); ¹³C NMR (DMSO-*d*₆/125 MHz): 124.69, 128.57, 130.73, 133.65, 139.26, 143.67, 148.32, 175.81; MS (ESI): 373 (M + H⁺); Anal. Calcd for C₁₅H₁₂N₆O₄S (372.36): C, 48.38; H, 3.25; N, 22.57; S, 8.61. Found: C, 48.21; H, 3.22; N, 22.65; S, 8.63.

Bis(3-nitrobenzaldehyde) thiocarbohydrazone (13)

Yellow powder (0.73 g, 98%): mp 222°C (dec.); IR (KBr): 3270, 3112, 1613, 1529, 1515, 1353, 1251, 876, 743, 677 cm⁻¹; ¹H NMR (DMSO-*d*₆/500 MHz): δ 7.77 (t, *J* = 8, 2H, aromatic H-3,3'), 8.28 (dd, *J* = 8, *J* = 1.5, 2H, aromatic H-2,2'), 8.28 (br s, 2H, H-4,4'), 8.66 (br s, 4H, aromatic H-1,1' and 2 azomethine H), 12.05 (br s, 1H, N-H), 12.16 (br s, 1H, N-H); ¹³C NMR (DMSO-*d*₆/125 MHz): 121.43, 124.23, 130.30, 133.56, 135.99, 141.00, 148.29, 175.43; MS (ESI): 373 (M + H⁺); Anal. Calcd for C₁₅H₁₂N₆O₄S (372.36): C, 48.38; H, 3.25; N, 22.57; S, 8.61. Found: C, 48.34; H, 3.24; N, 22.68; S, 8.60.

Bis(4-nitrobenzaldehyde) thiocarbohydrazone (14)

Yellow powder (0.58 g, 78%): mp 217°C (dec.); IR (KBr): 3284, 3153, 1545, 1523, 1377, 1327, 1260, 850 cm⁻¹; ¹H NMR (DMSO-*d*₆/500 MHz): δ 8.03-8.22 (m, 5H, 4 aromatic H and 1 azomethine H), 8.32 (d, *J* = 8.05, 4H, aromatic H-3,5,3',5'), 8.73 (br s, azomethine H), 12.02 (br s, N-H), 12.34 (br s, N-H); ¹³C NMR (DMSO-*d*₆/125 MHz): 123.86, 128.17, 140.31, 146.36, 147.80, 175.50; MS (ESI): 373 (M + H⁺); Anal. Calcd for C₁₅H₁₂N₆O₄S (372.36): C, 48.38; H, 3.25; N, 22.57; S, 8.61. Found: C, 48.18; H, 3.24; N, 22.65; S, 8.63.

Bis(pyridine-2-carbaldehyde) thiocarbohydrazone (15)

Yellow powder (0.49 g, 86%): mp 204-207°C; IR (KBr): 3140, 1612, 1538, 1524, 1249, 1126, 912, 886, 797 cm⁻¹; ¹H NMR (DMSO-*d*₆/500 MHz): (for the major isomer) δ 7.42 (ddd, *J* = 7.6, *J* = 4.5, *J* = 1.0, 2H, aromatic H-4,4'), 7.90 (dt, *J* = 7.6, *J* = 1.4, 2H, aromatic H-5,5'), 7.99 (br s, 1H, aromatic H-6), 8.22 (br s, 1H, aromatic H-6'), 8.37 (br s, 1H, azomethine H), 8.61 (d, *J* = 4.5, 2H, aromatic H-3,3'), 8.64 (br s, 1H, azomethine H), 11.90 (br s, 1H, N-H), 12.22 (br s, 1H, N-H); ¹³C NMR (DMSO-*d*₆/125 MHz): 119.65, 120.22, 124.44, 124.57, 125.05, 126.71, 136.75, 137.07, 138.56, 139.44, 143.88, 148.29, 149.55, 149.88, 151.37, 152.82, 175.49, 176.75; MS (ESI): 285 (M + H⁺), 307 (M + Na⁺); Anal. Calcd for C₁₃H₁₂N₆S (284.34): C, 54.91; H, 4.25; N, 29.56; S, 11.28. Found: C, 55.00; H, 4.23; N, 29.48; S, 11.26.

Bis(pyridine-3-carbaldehyde) thiocarbohydrazone (16)

Pale yellow powder (0.56 g, 98%): mp 215°C; IR (KBr): 3298, 3143, 1602, 1539, 1309, 1275, 1153, 821, 717 cm⁻¹; ¹H NMR (DMSO-*d*₆/500 MHz): δ 7.50 (dd, *J* = 7.64, *J* = 4.75, 2H, pyridine H-5,5'), 8.17 (br s, 2H, pyridine H-6,6'), 8.30 (br s, 1H, pyridine H-2), 8.62 (dd, *J* = 4.75, *J* = 1.46, pyridine H-4,4'), 8.66 (br s, 1H, pyridine H-2'), 8.66 (br s, 1H, azomethine H) 9.04 (br s, 1H, azomethine H), 11.78 (br s, 1H, N-H), 12.14 (br s, 1H, N-H); ¹³C NMR (DMSO-*d*₆/125 MHz): 123.78, 129.97, 133.77, 140.63, 146.05, 148.75, 150.53; MS (ESI): 285 (M + H⁺), 307 (M + Na⁺); Anal. Calcd for C₁₃H₁₂N₆S (284.34): C, 54.91; H, 4.25; N, 29.56; S, 11.28. Found: C, 54.68; H, 4.27; N, 29.53; S, 11.27.

Bis(pyridine-4-carbaldehyde) thiocarbohydrazone (17)

Yellow powder (0.56 g, 98%): mp 218-219°C; IR (KBr): 3214, 1597, 1562, 1511, 1261, 1131, 999, 907, 814 cm⁻¹; ¹H NMR (DMSO-*d*₆/500 MHz): δ 7.71 (br s, 2H, pyridine H-2,6), 7.82 (br s, 2H, pyridine H-2',6'), 8.14 (br s, 1H, azomethine H), 8.62 (br s, 1H, azomethine H), 8.67 (d, *J* = 5.85, 4H, H-3,5,3',5'), 11.94 (br s, 1H, N-H), 12.31 (br s, 1H, N-H); ¹³C NMR (DMSO-*d*₆/125 MHz): 121.24, 141.27, 146.84,

150.12, 175.65; MS (ESI): 307 (M + Na⁺); Anal. Calcd for C₁₃H₁₂N₆S (284.34): C, 54.91; H, 4.25; N, 29.56; S, 11.28. Found: C, 54.90; H, 4.23; N, 29.51; S, 11.23.

Bis(furan-2-carbaldehyde) thiocarbohydrazone (18)

Yellow powder (0.46 g, 88%): mp 186°C (dec.); IR (KBr): 3292, 3128, 1618, 1530, 1242, 1020, 944, 767, 624 cm⁻¹; ¹H NMR (DMSO-*d*₆/500 MHz): δ 6.6 (s, 2H, furan H-4,4'), 6.96 (br s, 2H, furan H-5,5'), 7.86 (s, 2H, furan H-3,3'), 8.05 (br s, 1H, azomethine H), 8.46 (br s, 1H, azomethine H), 11.46 (br s, 1H, N-H), 11.80 (br s, 1H, N-H); ¹³C NMR (DMSO-*d*₆/125 MHz): 112.81, 114.17, 134.50, 139.10, 145.69, 149.71, 174.88; MS (ESI): 285 (M + Na⁺); Anal. Calcd for C₁₁H₁₀N₄O₂S (262.29): C, 50.37; H, 3.84; N, 21.36; S, 12.23. Found: C, 50.47; H, 3.82; N, 21.27; S, 12.21.

Bis(thiophene-2-carbaldehyde) thiocarbohydrazone (19)

Yellow powder (0.41 g, 70%): mp 196°C (dec.); IR (KBr): 3294, 3114, 1589, 1543, 1526, 1258, 1224, 1130, 707 cm⁻¹; ¹H NMR (DMSO-*d*₆/500 MHz): δ 7.15 (dd, *J* = 4.86, *J* = 3.63, 2H, thiophene H-4,4'), 7.48 (br s, 2H, thiophene H-5,5'), 7.70 (d, *J* = 4.86, 2H, thiophene H-3,3'), 8.37 (br s, 1H, azomethine H), 8.68 (br s, 1H, azomethine H), 11.40 (br s, 1H, N-H), 11.76 (br s, 1H, N-H); ¹³C NMR (DMSO-*d*₆/125 MHz): 127.95, 129.22, 130.99, 138.51, 143.98, 173.97; MS (ESI): 295 (M + H⁺), 317 (M + Na⁺); Anal. Calcd for C₁₁H₁₀N₄S₃ (294.42): C, 45.87; H, 3.42; N, 19.03; S, 32.67. Found: C, 45.03; H, 3.40; N, 19.09; S, 32.76.

Bis(acetophenone) thiocarbohydrazone (20)

Recrystallized from ethanol; yellow powder (0.39 g, 63%): mp 188°C (dec.); IR (KBr): 3282, 3191, 1511, 1478, 1125, 791, 683 cm⁻¹; ¹H NMR (DMSO-*d*₆/500 MHz): δ 2.38 (s, 6H, CH₃), 7.45 (m, 6H, aromatic H-3,4,5,3',4',5'), 7.88 (d, *J* = 7.80, 4H, aromatic H-2,6,2',6'), 10.96 (br s, 2H, NH); MS (ESI): 311 (M + H⁺); Anal. Calcd for C₁₇H₁₈N₄S (310.42): C, 65.78; H, 5.84; N, 18.05; S, 10.33. Found: C, 65.60; H, 5.83; N, 18.01; S, 10.36.76.

Bis(N-(4-acetylphenyl) acetamide) thiocarbohydrazone (21)

Recrystallized from ethyl acetate; yellow powder (0.23 g, 54%): mp 220°C (dec.); IR (KBr): 3254, 3380, 3245, 1667, 1511, 1499, 1229, 850 cm⁻¹; ¹H NMR (DMSO-*d*₆/500 MHz): δ 2.06 (s, 6H, -C(O)CH₃), 2.34 (s, 6H, -C(N)CH₃), 7.64(d, *J* = 8.69, 4H, aromatic H-3,5,3',5'), 7.82 (d, *J* = 8.69, 4H, aromatic H-2,6,2',6'), 10.10 (s, 2H, NH), 10.76 (br s, 2H, NH); MS (ESI): 425 (M + H⁺); Anal. Calcd for C₂₁H₂₄N₆O₂S (424.52): C, 59.41; H, 5.70; N, 19.80; O, 7.54; S, 7.55. Found: C, 59.31; H, 5.71; N, 19.86; S, 32.73.

Bis(4'-fluoroacetophenone) thiocarbohydrazone (22)

Recrystallized from ethanol; yellow powder (0.37 g, 53%): mp 212-215°C; IR (KBr): 2365, 3150, 1595, 1487, 1225, 847, 834 cm⁻¹; ¹H NMR (DMSO-*d*₆/500 MHz): δ 2.36 (s, 6H, CH₃), 7.28 (t, *J* = 8.80, 4H, aromatic H-3,5,3',5'), 7.94 (dd, *J* = 8.80, *J* = 5.6, 4H, aromatic H-2,6,2',6'), 10.80 (br s, 2H, NH); MS (ESI): 347 (M + H⁺); Anal. Calcd for C₁₇H₁₆F₂N₄S (346.40): C, 58.94; H, 4.66; N, 16.17; S, 9.26. Found: C, 59.01; H, 4.65; N, 16.22; S, 9.25.

Bis(4'-chloroacetophenone) thiocarbohydrazone (23)

Recrystallized from ethanol; yellow powder (0.45 g, 59%): mp 213-216°C; IR (KBr): 3290, 1606, 1484, 1319, 1230, 1138, 830, 767 cm⁻¹; ¹H NMR (DMSO-*d*₆/500 MHz): δ 2.36 (s, 6H, CH₃), 7.51 (d, *J* = 8.61, 4H, aromatic H-3,5,3',5'), 7.91 (d, *J* = 8.61, 4H, aromatic H-2,6,2',6'), 10.87 (s, 2H, NH); MS (ESI): 379, 381, 383 (M + H⁺); Anal. Calcd for C₁₇H₁₆Cl₂N₄S (379.31): C, 53.83; H, 4.25; N, 14.77; S, 8.45. Found: C, 53.95; H, 4.25; N, 14.73; S, 8.47.

Bis(4'-bromoacetophenone) thiocarbohydrazone (24)

Yellow powder (0.85 g, 92%): mp 209°C (dec.); IR (KBr): 3290, 3183, 1596, 1509, 1477, 1229, 1008, 853 cm⁻¹; ¹H NMR (DMSO-*d*₆/500 MHz): δ 2.35 (s, 6H, CH₃), 7.64 (d, *J* = 8.50, 4H, aromatic H-3,5,3',5'), 7.83 (d, *J* = 8.50, 4H, aromatic H-2,6,2',6'), 10.88 (s, 2H, NH); MS (ESI): 467, 469, 471 (M + H⁺); Anal. Calcd for C₁₇H₁₆Br₂N₄S (468.21): C, 43.61; H, 3.44;

N, 11.97; S, 6.85. Found: C, 43.70; H, 3.43; N, 11.93; S, 6.84.

Bis (2-acetylpyridine) thiocarbohydrazone
(25)

Recrystallized twice from ethanol; yellow powder (0.32 g, 51%): mp 163-165°C; IR (KBr): 3192, 1583, 1551, 1463, 1227, 1118, 855 cm⁻¹; ¹H NMR (DMSO-*d*₆/500 MHz): δ 2.47 and 2.49 (each s, 3H, CH₃), 7.44 (dd, *J* = 7.4, *J* = 4.8, 2H, pyridine H-5,5'), 7.90 (t, *J* = 7.8, 2H, pyridine H-4,4'), 8.19 (d, *J* = 7.8, 2H, pyridine H-3,3'), 8.63 (d, *J* = 4.8, 2H, pyridine H-6,6'), 11.00 (br s, 2H, NH); MS (ESI): 313 (M + H⁺); Anal. Calcd for C₁₅H₁₆N₆S (312.39): C, 57.67; H, 5.16; N, 26.90; S, 10.26 Found: C, 57.54; H, 5.16; N, 26.96; S, 10.23.

Bis(4-acetylpyridine) thiocarbohydrazone
(26)

Yellow powder (0.40 g, 64%): mp 233-236°C; IR (KBr): 3312, 3147, 1593, 1510, 1486, 1217, 1111, 815cm⁻¹; ¹H NMR (DMSO-*d*₆/500 MHz): δ 2.39 (s, 6H, CH₃), 7.82 (d, *J* = 4.60, 4H, aromatic H-2,6,2',6'), 8.65 (d, *J* = 4.60, 4H, aromatic H-3,5,3',5'), 11.10 (s, 2H, NH); MS (ESI): 313 (M + H⁺); Anal. Calcd for C₁₅H₁₆N₆S (312.39): C, 57.67; H, 5.16; N, 26.90; S, 10.26. Found: C, 57.50; H, 5.18; N, 26.82; S, 10.25.

In-vitro evaluation of anti-mycobacterial activity

To evaluate the antimycobacterial activity of the compounds, the broth microdilution method (12) against *M. bovis* BCG (1173P2) was used. The test compounds were initially dissolved in DMSO to give a concentration of 1 or 2 mg/L. All wells of microplates received 100 μL of freshly prepared Middle broke 7H9 medium (Himedia, India), except the first column. Two hundred μL of distilled water was added to the first column of 96 well plates to minimize the evaporation of the medium in the test wells during the incubation. Then, 100 μL of test compounds with desired concentrations (1000 or 2000 μL) were added to the wells of the first row (each concentration was assayed in duplicate) and serial dilution was made from the first row to the last. Microbial suspension of BCG (1173P2) (100 μL), which had been prepared with standard

concentration of 0.5 Mcfarland and diluted with 1:10 proportion by the distilled water, was added to all test wells. Plates were then sealed and incubated for 4 days at 37°C. After that, 12 μL Tween 80 10% and 20 μL Alamar blue 0.01% (Himedia, India) were added to each test well. The results were assessed after 24 and 48 h. A blue color was interpreted as no bacterial growth, and color change to pink was scored as bacterial growth. Wells with a well-defined pink color were scored as positive for growth. The MIC (Minimal Inhibition Concentration) was defined as the lowest drug concentration, which prevented a color change from blue to pink. Ethambutol (Iranbaru, Tehran) and thiacetazone were used as positive control and DMSO as negative one.

In-vitro evaluation of antifungal activity

The antifungal activity was evaluated by the modified antimicrobial susceptibility testing based on NCCLS M27-A method (13) against *Candida albicans* ATCC 10231. The compounds were dissolved in dimethyl sulfoxide (DMSO) to reach the concentration of 0.5 or 1 mg/mL. The absorbance was read at 530 nm for fungi inoculums to reach the suitable density of microorganisms in order to yield the desired transmittance that is 75-77% equal to a particular fungal density. Working fungal culture was prepared from the stock fungal culture, 1:1000 dilution with broth (e.g. 10 μL stock fungal culture: 10 mL broth). Sabouraud maltose broth (DIFCO, Becton, Dickinson, USA) was used as the growth medium. Broth (100 μL) was added to each well of a 96-well microplate and then 40 μL of compounds and 60 μL broth were added to well (A), then a solution (100 μL) serially diluted from well (A) by taking 100 μL into (B) was obtained. This two-fold dilution was continued down the plate and 100 μL from the last well (H) was discarded. Then, all the wells were filled with 100 μL of working fungal culture. Itraconazole and Amphotericin B were used as a reference in the antifungal test. For this experiment and the following controls were prepared wells containing serial dilution of DMSO and itraconazole only. The plate was covered and incubated at 37°C for 24-48 h. The MIC values were obtained by reading the concentration of the well with no growth.




Figure 3. The route for the preparation of thiocarbohydrazones; reaction conditions: (i) H_2O , reflux. (ii) H_2O /Ethanol and acetic acid (catalytic amount), reflux.

Brine shrimp toxicity study

Brine shrimp lethality bioassay (14-17) was performed to investigate the toxicity of compounds 8, 12, 19 and 25 which showed the highest antimycobacterial activity. Dried cysts (1 g cyst per liter) of brine shrimp (*Artemia salina*) were hatched in a bottle containing artificial sea water (3.5% (w/v) marine salts/distilled water) at 28-30°C with strong aeration (flow rate of 7 L/min), under a continuous light regime (1600 lux) for 30-35 h. Subsequently, the newly hatched brine shrimp larvae (nauplii) were separated from the remaining cysts and collected with a pipette from the lighted side and concentrated in Petri dishes to be immediately used for bioassay. Assays were performed in 24-well flat test plates (Orange Scientifique, Belgium). Acetone 100% (Merck, Germany) was used for the preparation of different concentrations (1000, 100, 10 and 1 $\mu\text{g}/\text{mL}$) of tested compounds, in triplicates. Each well of treated groups became exposed to several concentration of acetone solution of compounds in the basic salt medium (3.5% (w/v) marine salts/distilled water in addition to polyethylene glycol (PEG) 6000 (Merck, Germany) 1.2%, while control groups only received basic salt medium. Gallic acid (Merck, Germany) was utilized as positive control. After the evaporation of vehicle solvent, 10 fresh nauplii were introduced to each well and the plates were placed on a shaker with 40 rpm to be aerated at room temperature. After 24 h, the numbers of survivors (larvae

were considered dead if they did not exhibit any internal or external movement during several seconds of observation) were counted by microscope AC 230V, 50 Hz (Sairan, Iran) and recorded to determine the corrected mortality via the following formula:

$$\text{Corrected mortality (\%)} = \frac{[(Mm_{ct_t}) - (Mm_{ct_c}) / 100 - (Mm_{ct_c})]}{\times 100}$$

Here: Mm_{ct} (mortality of individuals at time $t\%$) = $[\frac{N_{Mm}}{N_0}$ (number of died individuals) / N_0 (initial number of living individuals in every test well at the beginning of the test)] $\times 100$

(Mm_{ct_t}) = calculated Mm_{ct} for treated test wells

(Mm_{ct_c}) = calculated Mm_{ct} for control test wells

Based on the calculated corrected mortality, relevant 50% lethality doses (LD_{50})s with 95% confidence intervals were estimated by GraphPad Prism 5.0 (2007) for each tested compound.

In silico calculation of the physicochemical properties

The ClogP values were obtained through server tool available at (www.organic-chemistry.org). The other properties were measured by Toolkit for Estimating Physicochemical Properties of Organic Compounds (V. 1.0, 1999). The molecules in sdf format were made into database using EdiSDFd (V. 5.03, 2010). After




Figure 4. General structure for compounds 1, 3, 4, 5, 7, 8, 10, 11, 16, 18

that the energy minimization was performed through MMFF94 Energy Local Minimum, the group matching for the ranking based on key points in thiocetazone pharmacophores was measured in LigandScout (V.3.0, 2011). Dipole moment was calculated in Chem3D module of ChemBiooffice 12.0 package (2010). Energy minimization was first run by MM2, and then dipole was calculated by GAMESS interface.

Results and discussion

Chemistry

The overall synthetic route is summarized in Figure 3. Thiocarbohydrazide was the key intermediate which was readily synthesized based on the method reported in the literature (18) (30% yield). The next step was a Schiff base formation which was performed by the reaction between one mole equivalent of thiocarbohydrazide and two mole equivalents of different aromatic aldehydes or methyl ketones. This reaction was quite convenient with satisfying yields ranging from 55 to 90%.

The IR, $^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ spectra were consistent with the desired derivatives which were also confirmed by electrospray (ESI) mass spectrometry.

Several of the synthesized compounds (1, 3, 4, 5, 7, 8, 10, 11, 16, 18) showed two olefinic carbons in $^{13}\text{C-NMR}$ spectra suggesting the existence of two diastereomeric carbazone moieties in these compounds (Figure 4).

Compounds 2, 6, 9, 12, 13, 14, 17 and 19 on the other hand showed symmetric structures and no diastereomeric moieties were recognized in their $^{13}\text{C-NMR}$ spectra. Only compound 15 showed the existence of the mixture of different isomers a-c (Figure 5).

$^1\text{H-NMR}$ spectrum of this compound shows two series of peaks for the pyridine ring confirming the existence of both a and b isomers.




Figure 5. Compound 15 as different isomers.

Comparing the integration of these two series of peaks suggests the existence of 2:1 ratio for the isomers. Besides, these two series of peaks, the presence of one extra triplet and one extra doublet peak in $^1\text{H-NMR}$ suggest the existence of c isomer in the mixture. Attempts to separate different isomers of compound 15 were not successful probably due to the interconversion of the isomers during the purification process.

Another plausible explanation for the above mentioned extra olefinic hydrogen and carbon signals in $^1\text{H NMR}$ and $^{13}\text{C NMR}$ spectra could be the possibility of the formation of intramolecular hydrogen bond between the imine nitrogen and thioamide N-H (Figure 6). This intramolecular H-bond is possible in both Z and E isomers of thiocarbazone compounds. Involvement of the imine nitrogen in the intramolecular H-bond will cause a difference in electronic environment of the two olefinic hydrogens and carbons in thiocarbazone compounds and therefore they appear at different chemical shifts. The existence of intramolecular H-bond of this type has been proven by X-ray crystallography in crystals of thiosemicarbazone derivatives before (19, 20).




Figure 6. Existence of possible intramolecular H-bond in both E and Z isomers. H-bonds are shown by dashed lines.




Figure 7. Proposed symmetric structure for compounds 20-26 based on ^1H NMR spectra.

^1H NMR spectra obtained for the synthesized derivatives in the present study suggest the existence of the same type of H-bonding in dimethylsulfoxide as the solvent.


However, in the case of the thiosemicarbazones of methyl ketones (20-26), ^1H NMR data suggests a symmetric structure in which the C=S bond is located on the symmetry line of the molecule (Figure 7). In the ^1H NMR spectra of these derivatives, each proton in one side of the symmetry line is the chemical shift equivalent of the corresponding proton in the other side. This could be for the reason that the bulky methyl groups limit the flexibility of the molecule needed to adopt the conformation suitable for intramolecular hydrogen bonding as shown in Figure 6 and the molecule seems to prefer a symmetric conformation.

Biological activity

The *in-vitro* antimycobacterial and antifungal assays were performed at Pasteur Institute (Tehran, Iran). The antimycobacterial activity of the compounds was measured by the broth microdilution method (12) against *M. bovis* BCG (1173P2) and ethambutol and thiacetazone were used as standard controls. The *in-vitro* antifungal activity evaluation was performed by the modified antimicrobial susceptibility testing based on NCCLS M27-A method (13). *Candida albicans* ATCC 10231 was used as test strain and itraconazole and amphotericin B as standard controls. The MIC (Minimal Inhibition Concentration) was defined as the lowest concentration which could inhibit the mycobacterial or fungal growth. The detailed procedure is described in the experimental section.

The synthesized derivatives and their antimycobacterial activities are listed in Table

1. The most active compounds are 25 (MIC < 1.9 $\mu\text{g}/\text{mL}$), 19 and 8 with MICs of 15.6 $\mu\text{g}/\text{mL}$. The MICs for standard controls (thiacetazone and ethambutol) were obtained less than 3.9 and 0.75 $\mu\text{g}/\text{mL}$, respectively. Compounds 11, 12, 15 and 18 had acceptable activities (MIC = 31.3 $\mu\text{g}/\text{mL}$) among the other analogs. Calculation of general physicochemical properties for compounds 1-26 *i.e.* ClogP, dipole moment, and logP (O/W) are shown in Table 2. Attempts to correlate these properties to the activity of the compounds reveal some relationship. Barry *et al.* (21) have identified that the optimum logP range for antimycobacterial property is 1.3-4.1. The calculated logP values indicated in Table 2 shows that compounds 15, 18, 19 and 25 with strong activity fall into this category. However, as identified in that study, lipophilic character alone in regard to the molecular bioactivity may not correlate well with increased antimicrobial activity. On the other hand, *in-vitro* studies of thiosemicarbazone analogs perhaps offer the clearest demonstration of activity versus optimal hydrophobicity (22). For example, this study afforded the correlation of logP and the type of *Mycobacterium* tested in the bioassay. They indicated that for *M. tuberculosis* and a variety of slow-growing mycobacteria, the optimum logP value was 4, whereas for a fast-growing strain such as *M. smegmatis*, the optimum was 3. LogP in higher range contributes to a shift from optimum needed values that may possibly prevent the compound to reach the biological target points due to absorption to the cellular lipids. Many physicochemical factors can in a complicated manner affect the activity of antimycobacterial agents (23). Dipole moment is among the factors that influence the activity of the compounds. In general, the lower the dipole moment, the better would be the antimycobacterial activity. This

Table 1. Antimycobacterial activity data of the synthesized compounds.

Derivative	Ar	R	MIC ($\mu\text{g/mL}$)	
			24 h	48 h
1	Phenyl	H	125	125
2	4-acetamidophenyl	H	125	187.5
3	2-fluorophenyl	H	125	125
4	3-fluorophenyl	H	125	125
5	4-fluorophenyl	H	125	125
6	2-chlorophenyl	H	125	125
7	3-chlorophenyl	H	46.8	62.5
8	4-chlorophenyl	H	15.6	15.6
9	2-bromophenyl	H	93.7	125
10	3-bromophenyl	H	62.5	62.5
11	4-bromophenyl	H	31.2	31.2
12	2-nitrophenyl	H	31.2	31.2
13	3-nitrophenyl	H	187.5	250
14	4-nitrophenyl	H	187.5	250
15	2-pyridyl	H	31.2	46.8
16	3-pyridyl	H	125	125
17	4-pyridyl	H	62.5	125
18	2-furyl	H	31.2	31.2
19	2-thienyl	H	15.6	15.6
20	Phenyl	CH ₃	125	125
21	4-acetamidophenyl	CH ₃	125	187.5
22	4-fluorophenyl	CH ₃	125	125
23	4-chlorophenyl	CH ₃	125	125
24	4-bromophenyl	CH ₃	125	125
25	2-pyridyl	CH ₃	< 1.9	< 1.9
26	4-pyridyl	CH ₃	125	125
Ethambutol			0.75	0.75
Thiacetazone			< 3.9	< 3.9
DMSO			6.5% v/v	6.5% v/v

is particularly indicated in compounds 12 and 15. However, a general weak activity is seen in fluorinated derivatives, while in the chlorinated and brominated compounds, the bioactivity is mainly seen in 4-substituted ones in accordance with the pharmacophore score increase. Among these, low dipole value is seen among the less active compounds that matches the pattern of increase in pharmacophore score. Our results for the possible effect of the minimized shape

of molecules compared to thiacetazone provide certain affirmative clues on such correlations. In this regard, compounds 8, 11, 19 and 25 show high spatial pharmacophore arrangement after the alignment with thiacetazone (Table 2). Although there are very high pharmacophore score values for compounds 2 and 21, as the highest, these compounds do not seem to present other suitable values, such as lipophilicity ($\log P$ (O/W)) and in particular dipole moment, in the

Table 2. The predicted properties and the pharmacophore fitness score data of synthesized compounds.

Compound	Dipole (Debye)	logP (O/W) ¹	ClogP ²	Pharmacophore score ^{3,5}
1	6.77	3.00	3.72	68.57
2	6.06	1.84	2.93	88.62
3	5.51	3.28	3.84	37.54
4	4.03	3.28	3.84	58.15
5	6.07	3.28	3.84	68.67
6	5.41	4.24	4.95	57.97
7	4.33	4.24	4.95	68.62
8	6.12	4.24	4.95	68.68
9	4.65	4.78	5.12	68.44
10	4.39	4.78	5.12	57.98
11	6.23	4.78	5.12	68.68
12	4.54	2.92	3.46	57.53
13	2.43	2.92	3.46	68.22
14	7.64	2.92	3.46	67.92
15	5.52	1.25	1.57	37.09
16	4.28	0.78	1.57	56.79
17	5.91	0.78	1.57	68.75
18	7.42	3.28 ⁴	1.93	57.98
19	7.55	2.74 ⁴	3.41	68.75
20	7.13	2.92	4.19	68.66
21	7.82	1.75	3.40	88.17
22	7.44	3.19	4.31	68.73
23	7.07	4.16	5.41	68.72
24	7.02	4.70	5.58	68.74
25	8.51	1.17	2.24	68.48
26	6.97	0.70	2.03	68.61
Thiacetazone	8.53	0.45	0.90	89.00

¹Obtained by Toolkit application; ²obtained from organic-chemistry.org server; ³measured based on thiacetazone; ⁴measured by GCM of Viswanadhan method; ⁵Since the data in this column is calculated based on thiacetazone structural features, the data is only used to study the structure-antitubercular activity relationship related to antimycobacterial property of the compounds.

optimum needed range.

Thiacetazone is a second-line drug for the treatment of tuberculosis. From the chemical standpoint, thiacetazone belongs to thiocarbonyl derivatives and it has long been known that these derivatives could be prodrugs that have to be converted to their active forms by mycobacterial enzymes (24). In 2000, two independent research groups reported that EthA is involved in the activation of thiocarbonyl drugs in both *M. tuberculosis* and *M. leprae* and resistance of *M. tuberculosis* to thiacetazone involves mutations in EthA (20, 22). These findings suggest that

compounds 1-26 could also be considered as prodrugs which need to be converted to their active S-oxide form by a flavin containing monooxygenase in mycobacteria. The active forms will then interfere with some essential and vital processes in *Mycobacterium* cell such as cyclopropanation of mycolic acid (25).

Considering the fact that antimycobacterial effect of thiocarbohydrazone derivatives 1-26 depends on two independent processes, *i.e.* metabolic activation and inhibition of cyclopropanation of mycolic acid in *Mycobacterium*, it is not surprising that no




Figure 8. Metabolic activation of thiacetazone by a flavin-containing monooxygenase in *Mycobacterium*.

decisive structure-activity relationship could be concluded for these compounds.

Besides the antitubercular activity, the antifungal activity of the new derivatives against *C. albicans* is noticeable. Compound 25 exhibited the highest activity (MIC < 3.25 µg/mL) against *C. albicans*. The derivatives with acceptable activity include compounds 8 and 15 (MIC = 15.63 µg/mL) and 19 (MIC = 62.5 µg/mL) (Table 3). Other derivatives showed weak (if any) activity. The important point is that compounds which are active against *C. albicans* have excellent or acceptable antitubercular activity as well. In this regard, compound 25 is superior to all, since it is the most active compound against both tested organisms. This finding is valuable in order to conduct the mechanistic studies in the future on the targets that play vital roles in both organisms and may lead to revealing less noticed metabolic pathways, in which the newly synthesized derivatives may interfere.

Determining the cytotoxicity properties of given drug candidates is crucial to their fate in lead identification and the following phases of drug discovery process. In recent years, there has been a number of toxicity tests developed in which the response has been demonstrated in invertebrates. Among these tests, the brine shrimp lethality bioassay (14, 15, 16 and 26) enjoys the advantages of being inexpensive, reproducible, conveniently handled and environmentally relevant which make it a practical method for preliminary assessment of toxicity. By carefully controlling the factors such as temperature, composition, salinity of the medium and the age of the larvae, the obtained result will be accompanied by satisfactory repeatability. Though insufficient to deduce the mechanism of action, the brine shrimp bioassay is vastly used as primary toxicity assessment of natural and synthetic leads (15). It has been demonstrated that the nature of the systems in

Table 3. Antifungal activity of the synthesized compounds against *C. albicans* using broth dilution method.

Derivative	MIC (µg/mL)	
	(24 h)	(48 h)
1	500	1000
2	1000	> 1000
3	500	1000
4	500	1000
5	1000	> 1000
6	125	500
7	500	1000
8	15.625	31.25
9	1000	1000
10	250	1000
11	62.25	125
12	1000	1000
13	500	1000
14	500	1000
15	15.625	31.25
16	500	1000
17	1000	1000
18	250	500
19	62.5	125
20	500	1000
21	500	1000
22	500	1000
23	500	1000
24	1000	1000
25	< 3.25	< 3.25
26	500	1000
Itraconazol	0.5	2
Amphotericin B	< 0.25	0.25
DMSO	10% v/v	10% v/v




Figure 9. Non-linear regression curve for brine shrimp toxicity of compounds 8, 12, 19 and 25.

brine shrimp which responds to drugs appears to be similar to the mammalian systems. As a result, this bioassay has been suggested for screening biological activities.

In this study, the acute toxicity of four of the synthesized derivatives with highest antimycobacterial potency in addition to considering their diversity, were investigated by means of the *A. salina* short-term bioassay at Pasteur Institute (Tehran, Iran). The related LD_{50} s (Figures 9) were calculated based on analytical method non-linear regression (dose response inhibition). The analyzed data demonstrated that the selected derivatives caused lethality to

nauplii with LD_{50} values ranging from 2.69 to 188.10 $\mu\text{g/mL}$ (Table 4).

To explore the selectivity of antimycobacterial potency for each tested compound, a selectivity index (SI) was determined by dividing LD_{50} to MIC. The range of SIs thus calculated was between 0.17 to 12.04. Based on given findings, it may indeed be true that tested compounds 8, 12 and 25 with SIs 12.04, 1.71 and 4.32 have discriminating toxicity against *Mycobacterium bovis BCG* in comparison to eukaryotic organism. There would be no doubt that considering SIs for candidate compounds can lead to presenting a set of new antimycobacterial candidates

Table 4. Data obtained by *Artemia* lethality bioassay.

Derivative	LD_{50} ($\mu\text{g/mL}$)	LD_{50} (95% Confidence Intervals)	SI
8	188.10	12.90 to 2741.00	12.04
12	53.22	16.07 to 176.20	1.71
19	2.69	0.06 to 122.80	0.17
25	8.23	2.94 to 23.01	4.32
Gallic acid	23.84	9.78 to 58.13	

with minor undesirable side effects for further investigations. In this study, data suggest that compound 25 is a potent anti-mycobacterial and antifungal compound with excellent MICs and acceptable selectivity index which shows its safety on the tested eukaryotic organism. Further focused analog synthesis and antitubercular/antifungal activity studies are underway for compounds 8, 19 and 25 which may lead to new derivatives with enhanced efficiency against *M. tuberculosis* and fungal species.

Conclusion

Amongst the 26 synthesized compounds, some of them exhibited a pronounced activity against *M. bovis* BCG and *C. albicans* with MIC values comparable with that of the standard drugs. These derivatives can be considered as starting points for further modifications to reach compounds with promising anti-tubercular and antifungal activity to enter into other complicated assays involving mechanism of action, *in-vivo* tests and perhaps clinical trials.

Acknowledgment

Authors wish to acknowledge and thank Pasteur Institute of Iran and Ministry of Health and Medical Education in Iran for partly funding of this project as grant number 456.

References

- (1) World Health Organization. Global tuberculosis control: epidemiology, strategy, financing. WHO report (2009). Available online at: http://www.who.int/tb/publications/global_report/2009/en/index.html.
- (2) Dorman SE and Chaisson RE. From magic bullets back to the magic mountain: the rise of extensively drug-resistant tuberculosis. *Nat. Med.* (2007) 13: 295-298.
- (3) Centers for Disease Control and Prevention (CDC). *MMWR Morb Mortal Wkly Rep.* (2006) 55: 301-305.
- (4) Kobarfard F. Tuberculosis and traditional medicine: fighting the oldest infectious disease using the oldest source of medicines. *Iranian J. Pharm. Res.* (2004) 2: 71-72.
- (5) Davidson PT and Le HQ. Drug treatment of tuberculosis. *Drugs* (1992) 43: 651-673.
- (6) Bermudez LE, Reynolds R, Kolonoski P, Aralar P, Inderlied CB and Young LS. Thiosemicarbazole (thiacetazone-like) compound with activity against *Mycobacterium avium* in mice. *Antimicrob. Agents. Chemother.* (2003) 47: 2685-2687.
- (7) Kobarfard F and Kauffman JM. Halogenated antituberculosis agents. United States patent (2003) US2003114531.
- (8) Dover LG, Alahari A, Gratraud P, Gomes JM and Bhowruth VE. EthA, a common activator of thiocarbamide-containing drugs acting on different mycobacterial targets. *Antimicrob. Agents. Chemother.* (2007) 51: 1055-1063.
- (9) Qian L and Ortiz de Montellano PR. Oxidative activation of thiacetazone by the *Mycobacterium tuberculosis* flavin monooxygenase EtaA and human FMO1 and FMO3. *Chem. Res. Toxicol.* (2006) 19: 443-449.
- (10) DeBarber AE, Mdluli K, Bosman M, Bekker LG and Barry CE. Ethionamide activation and sensitivity in multidrug resistant *Mycobacterium tuberculosis*. *Proc. Natl. Acad. Sci. USA* (2000) 97: 9677-9682.
- (11) Sardari S, Shaghghi B and Borna H. in Mendez-Vilas, A. Ed. Microorganisms in Industry and Environment, from scientific and industrial research to consumer products. *World Scientific, New Jersey*, (2011) 665.
- (12) Camacho-Corona MR, Ramirez-Cabrera MA, Santiago OG, Garza-Gonzalez E, Palacios Idel P and Luna-Herrera J. Activity against drug resistant-tuberculosis strains of plants used in Mexican traditional medicine to treat tuberculosis and other respiratory diseases. *Phytother. Res.* (2008) 22: 82-85.
- (13) Cuenca-Estrella M, Lee-Yang W, Ciblak MA, Arthington-Skaggs BA, Mellado E and Warnock DW. Comparative evaluation of NCCLS M27-A and EUCAST broth microdilution procedures for antifungal susceptibility testing of *Candida* species. *Antimicrob. Agents. Chemother.* (2002) 51: 3644-3647.
- (14) Borowitz JL and McLaughlin JL. Evidence for Calcium Channels in Brine Shrimp: Oiltlazem Protects Shrimp against Cadmium. *Bull. Environ. Contam. Toxicol.* (1992) 48:435-440.
- (15) Hartl M and Humpf HU. Toxicity assessment of fumonisins using the brine shrimp (*Artemia salina*) bioassay. *Food. Chem. Toxicol.* (2000) 38: 1097-1102.
- (16) Favilla M, Macchia L, Gallo A and Altomare C. Toxicity assessment of metabolites of fungal biocontrol agents using two different (*Artemia salina* and *Daphnia magna*) invertebrate bioassays. *Food. Chem. Toxicol.* (2006) 44: 1922-1931.
- (17) Baravalia Y, Vaghasiya Y and Chanda S. Brine Shrimp cytotoxicity, anti-inflammatory and analgesic properties of *Woodfordia fruticosa* Kurz Flowers. *Iranian J. Pharm. Res.* (2012) 11: 851-861.
- (18) Burns GR. Metal complexes of thiocarbonylhydrazide. *Inorg. Chem.* (1968) 7: 277-283.
- (19) Richardson DR, Kalinowski DS, Richardson V, Sharpe PC, Lovejoy DB, Islam M and Bernhardt PV. 2-Acetylpyridine thiosemicarbazones are potent iron chelators and antiproliferative agents: redox activity, iron complexation and characterization of their antitumor activity. *J. Med. Chem.* (2009) 52: 1459-1470.

- (20) Kalinowski DS, Yu Y, Sharpe PC, Islam M, Liao Y, Lovejoy DB, Kumar N, Bernhardt PV and Richardson DR. Design, synthesis, and characterization of novel iron chelators: structure–activity relationships of the 2-benzoylpyridine thiosemicarbazone series and their 3-nitrobenzoyl analogues as potent antitumor agents. *J. Med. Chem.* (2007) 50: 3716-3729.
- (21) Barry CE, Slayden RA, Sampson AE and Lee RE. Use of genomics and combinatorial chemistry in the development of new antimycobacterial drugs. *Biochem. Pharmacol.* (2000) 59: 221-231.
- (22) Morrison NE and Collins FM. Antimycobacterial activity of 2-acetylpyridine thiosemicarbazones in relation to their antileprosy activity. *Int. J. Lepr.* (1981) 49:180-186.
- (23) Bartzatt R, Cirillo SLG and Cirillo JD. Determination of the molecular properties effectuating the growth inhibition of *Mycobacterium tuberculosis* by various small molecule hydrazides. *Lett. Drug. Des. Disc.* (2008) 5: 162-168.
- (24) Baulard AR., Betts JC, Engohang-Ndong J, Quan S, McAdam RA, Brennan PJ, Loch C and Besra GS. Activation of the pro-drug ethionamide is regulated in mycobacteria. *J. Biol. Chem.* (2000) 275: 28326-28331.
- (25) Alahari A, Trivelli X, Guerardel Y, Dover LG, Besra GS, Sachhettini JC, Reynolds RC, Coxon GD and Kremer L. Thiacetazone, an antitubercular drug that inhibits cyclopropanation of cell wall mycolic acids in mycobacteria. *Plos one* (2007) 2: e1343.
- (26) Ferrão-Filho SA and Kozlowsky-Suzuki B. Cyanotoxins: Bioaccumulation and Effects on Aquatic Animals. *Mar. Drugs* (2011) 9: 2729-2772.
-

This article is available online at <http://www.ijpr.ir>