Original Article

Propofol Attenuates Toxic Oxidative Stress by CCl₄ in Liver Mitochondria and Blood in Rat

Akram Ranjbar^{a*}, Mohammad Sharifzadeh^b, Jamshid Karimi^c, Heidar Tavilani^c, Maryam Baeeri^b, Tavakol Heidary shayesteh^a and Mohammad Abdollahi^b

^a Department of Toxicology and Pharmacology, School of Pharmacy, Hamadan University of Medical Sciences, Hamadan, Iran.^b Faculty of Pharmacy, and Pharmaceutical Sciences Research Center, Tehran University of Medical Sciences, Tehran, Iran.^c Department of Biochemistry, School of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran.

Abstract

Anti-oxidant effects of propofol (2, 6-diisopropylphenol) were evaluated agains carbon tetrachloridet CCl_4 -induced oxidative stress in rat liver. 30 male rats were equally divided in to 6 groups (5 rats each). Group I (control), while Group II was given CCl_4 (3 mL/Kg/day, IP). Animals of Groups III received only propofol (10 mg/Kg/day, IP). Group IV was given propofol+ CCl_4 . Group V was administered vitamin E (alpha-tocopherol acetate 15 mg/Kg/day, SC) .Animals of Group VII received alpha-tocopherol acetate + CCl_4 once daily for two weeks. After treatment, blood and liver mitochondria were isolated. Anti-oxidant enzymes activity such as glutathione peroxidase (GPx), superoxide dismutase (SOD) and oxidative stress marker such as reduced glutathione (GSH) and lipid peroxidation (LPO) concentration were measured. Oxidative stress induced with CCl_4 in liver mitochondria was evident by a significant increase in enzymatic activities of GPx, SOD, and LPO and decreased of GSH and vailability of mitochondria. Propofol and vitamin E restored CCl_4 -induced changes in GSH, GPx, SOD and LPO in blood and liver mitochondria. CCl_4 decreased viability of mitochondria that was recovered by propofol and vitamin E. It is concluded that oxidative damage is the mechanism of toxicity of CCl_4 in the mitochondria that can be recovered by propofol comparable to vitamin E.

Keywords: Propofol; Liver mitochondria; Oxidative stress; Rat; CCl₄.

Introduction

Propofol (2, 6-diisopropylphenol) is a widely used intravenous sedative-hypnotic agent for both induction/maintenance of anesthesia and sedation of critically ill patients (1) Advantages of this agent over others of similar applications include lower incidence of side effects (2) and better quality of anesthesia (3). Propofol>s structure contains a phenolic hydroxyl group and thus resembles that of a-tocopherol (vitamin E), a natural anti-oxidant. As shown by both *in-vitro* and *in-vivo* studies, the anti-oxidant activity of propofol results partly from this phenolic chemical structure (4). Numerous studies have demonstrated anti-oxidant effects of propofol *in-vitro* (5, 6) and *in-vivo* (7), but actions of propofol on different cells are varied and multiple mechanisms may be involved (8). Propofol has been demonstrated to prevent oxidative stress-mediated endothelial cell activation and dysfunction induced by hydrogen peroxide (8, 9) and tumour necrosis factor-a (TNF-a) (10,

^{*} Corresponding author:

E-mail: akranjbar1389@yahoo.com

11). The anti-oxidant status of a cell determines its susceptibility to oxidative damage, and is usually altered in response to oxidative stress (12). When reactive oxygen species (ROS) generation overwhelms the anti-oxidant defense, the free radicals can interact with endogenous macromolecules and alter cellular function (12). The mitochondrial respiratory chain is the major source of intracellular generation of ROS and at the same time, an important target for the damaging effects of ROS (13).

Carbon tetrachloride (CCl_4) is known to induce reactive free radicals and induction of cell damage through covalent binding to the membrane proteins (14).

 CCl_4 is converted in to trichloromethyl radical (CCl_3) and its derivative peroxy trichloromethyl radical ($OOCCl_3$) by cytochrome P450 in liver microsomes. These free radicals are highly reactive and are capable of reacting with polyunsaturated fatty acids of the membranous system leading to oxidative injuries such as lipid peroxidation (15). The aims of this study were to investigate the anti-oxidative effects of propofol on liver mitochondrial function in rat treated with CCl_4 .

Experimental

Chemicals

Tetraethoxypropane (MDA), 2-thiobarbituric acid (TBA), trichloroace tic acid (TCA), n-butanol, propofol , sucrose, ethylenediamine tetra acetic acid (EDTA), Comassie blue, bovine serum album in (BSA), 4,5(dimethylthiazol-2yl)-2,5-diphenyltetrazolium bromide (MTT), GPx and SOD (Ransel kit, Randox Laboratories Ltd, Crumlin, UK), bioxytech GSH kit (Oxis Research, USA), Comassie blue, bovine serum albumin (BSA), 4,5-(dimethylthiazol-2-yl)-2,5diphenyltetrazolium bromide (MTT), were used in this study. All other chemicals were obtained from the Sigma.

Animals and treatments

Adult male Wistar rats weighing 180-250 g maintained on a 12-hour light/dark cycle with free access to tap water and standard laboratory chow were used. Animals were randomly divided into six groups (n = 5) and treated for

2 week intraperitoneally (IP). The groups were as follows: control group, propofol group, CCl_4 group, propofol and CCl_4 group, vitamin E group, and CCl_4 and vitamin E group.

Propofol was administered (10 mg/Kg/day, IP) alone or in combination with CCl_4 (0.2 mL/Kg /day, IP) and vitamin E as (15 mg/Kg/day, SC) One group of animals received only normal saline and was assigned as control. At the end of the treatment, 24 hours post the last dose of treatment, animals were killed, liver tissue was separated and stored in liquid nitrogen. Blood samples were collected from heart in heparinized tubes and plasma was isolated. The experiments were conducted according to the ethical rules approved by Institutional Review Board (IRB).

Preparation of liver mitochondria

The liver was removed and minced with small scissors in a cold manitol solution containing 0.225 M D-manitol, 75 mM sucrose, and 0.2 mM ethylenediaminetetraacetic acid (EDTA). The minced liver (30 g) was gently homogenized in a glass homogenizer with a Teflon pestle and then centrifuged at $700 \times g$ for 10 min at 4 °C at remove nuclei, unbroken cells, and other non-subcellular tissue. The supernatants were centrifuged at $7,000 \times g$ for 20 min. These second supernatants were pooled as the crude microsomal fraction and the pale loose upper layer, which was rich in swollen or broken mitochondria, lysosomes, and some microsomes, of sediments was washed away.

The dark packed lower layer (heavy mitochondrial fraction) was resuspended in the manitol solution and recentrifuged twice at $7,000 \times g$ for 20 min. The heavy mitochondrial sediments were suspended in Tris solution containing 0.05 M Tris-HCl buffer (pH 7.4), 0.25 M sucrose, 20 mM KCl, 2.0 mM MgCl₂, and 1.0 mM Na₂HPO₄ at 4 °C before assay (16).

Measurement of Cu/Zn-SOD

The activity of Cu/Zn-SOD was measured using a commercial kit (Ransod kit, Randox Laboratories Ltd, Crumlin, UK). Measurement of the enzyme was based on the generation of superoxide radicals produced by xanthine and xanthine oxidase and reacted with 2-(4-iodophenyl)-3-(4-nitrofenol) 5-phenyltetrazolium chloride (INT) to form a red formazan dye. The formazan was read at 505 nm. One unit of Cu/Zn- SOD was defined as the amount of enzyme necessary to produce 50% inhibition in the INT reduction rate.

Measurement of GPx

The amount of GPx was determined using a commercially available kit (Ransel kit, Randox Laboratories Ltd, Crumlin, UK) by measuring the rate of oxidation of NADPH at 340 nm. A unit of enzyme was expressed as the amount of enzyme needed to oxidize 1 nmol of NADPH oxidase/minute.

Measurement of reduced glutathione assay (GSH)

Level of reduced glutathione (GSH) was measured using colorimetric assay kit. The kit uses 5, 50-dithiobis-2-nitrobenzoic acid (DTNB) and glutathione reductase. The procedure was followed according to manufacturer's instruction and the levels were quantitated as micromolar GSH based on standard supplied along with the kit.

Measurement of lipid peroxidation (LPO)

The LPO product in tissues was determined by thiobarbituric acid reagent expressed as the extent of malondialdehyde (MDA) productions during an acid heating reaction. Briefly, the samples were diluted by 1.5 mL TCA (20% w/v) was added to 250 μ L o f this samples and centrifuged in 3000 g for 10 min. Then, the precipitation was dissolved in sulfuric acid and 1.5 mL of the mixture was added to 1.5 mL of TBA (0.2% w/v). The mixture was then incubated for 1 h in a boiling water bath. Following incubation, 2 m l of n-butanol was added, the solution centrifuged, cooled and the absorption of the supernatant was recorded in 532 nm. The calibration curve of tetraethoxypropane standard solutions was used to determine the concentrations of TBA+MDA adducts in samples (17).

Total Protein

The protein content was quantified by the method of Bradford. Concentrated Coomassie blue (G250) was diluted in 250 µL distilled

water, and then 750 μ L of this diluted dye was added to 50 μ L of sample. The mixture was incubated at room temperature for 10 min and an absorbance measurement was taken at 595 nm by a spectrophotometer. A standard curve was constructed by using bovine serum albumin ranging between 0.25 and 1 mg/mL (18).

Mitochondrial viability

This assay is a quantitative colorimetric method to determine mitochondrial toxicity or viability of liver mitochondria. It utilizes the yellow tetrazolium salt (MTT), which is metabolized by mitochondrial dehydrogenase enzyme from viable cells to yield a purple formazan reaction product that was determined spectrophotometrically at wavelength of 570 nm. The percentage (of control) of mitochondrial viability of each test sample was calculated (19).

Statistical analysis

Mean and standard error values were determined for all the parameters and the results were expressed as Mean + SEM. All data were analyzed with SPSS Version 18 employing one-way ANOVA followed by Tukey post hoc test. Duplicate of experiments were performed. Differences between groups was considered significant when P < 0.05.

Results

Lipid peroxidation (LPO)

In blood: CCl_4 caused a significant increase in LPO when compared to control (p-value < 0.001). Propofol caused a significant decrease in LPO when compared to CCl_4 group (p-value < 0.001). Vitamin E reduced LPO when compared to CCl_4 (p-value < 0.001). Co administration of Vitamin E with CCl_4 significantly reduced CCl_4 induced LPO (p-value < 0.001).

In liver mitochondria: CCl_4 caused a significant increase in LPO when compared to control (p-value < 0.001). Propofol caused a significant decrease in LPO when compared to CCl_4 group (p-value < 0.001). Vitamin E reduced LPO when compared to CCl_4 (p-value < 0.001). Coadministration of propofol or Vitamin E with CCl_4 significantly reduced CCl_4 induced



Figure 1. Lipid peroxidation (LPO) in plasma and liver mitochondria of rats. Values are the mean \pm SE (n = 5) ^{aa}Significantly different from control group at p < .05. ^{bb}Significantly different from CCl₄ group at p < .05. PF, Propofol; AT, (alpha-tocopherol; vitamin E). Duplicate bars from left to right represent plasma and liver mitochondria, respectively.

LPO (p-value = 0.050 and p-value = 0.010, respectively); Figure 1.

Superoxide dismutase

In blood: CCl_4 caused a significant increase in SOD activity when compared to control (p-value < 0.001). Propofol caused a significant decrease in SOD activity when compared to CCl_4 group (p-value < 0.001). Vitamin E reduced SOD activity when compared to CCl_4 (p-value < 0.001).

In liver mitochondria: Vitamin E reduced SOD activity when compared to CCl_4 (p-value = 0.014); Figure 2.

Glutathione peroxidase

In blood: Vitamin E reduced GPx activity when compared to CCL_4 (p-value = 0.004). Coadministration of propofol with Vitamin E significantly reduced CCl_4 induced SOD activity (p-value=0.003 and p-value=0.013, respectively).

In liver mitochondria: Vitamin E reduced GPx activity when compared to CCl_4 (p-value = 0.001); Figure 3.

Reduced glutathione

In plasma: Treatment with CCl_4 decreased GSH as compared to control (p-value = 0.002). Propofol increased GSH as compared to CCl_4 group (p-value = 0.002). Vitamin E induced GSH when compared to CCl_4 (p-value = 0.001).

In liver mitochondria: Administration of CCl_4 decreased GSH in comparison to controls (p-value < 0.001). Treatment with propofol increased GSH as compared to CCl_4 group (p-value < 0.001). Vitamin E significantly increased GSH level as compared to CCl_4 group (p-value <



Figure 2. Superoxide dismutase (SOD) activity in plasma and liver mitochondria of rats. Values are the mean \pm SE (n = 5) ^{as}Significantly different from control group at p < .05. ^{bb}Significantly different from CCl4 group at p < .05. PF, Propofol; AT, (alpha-tocopherol; vitamin E). Duplicate bars from left to right represent plasma and liver mitochondria, respectively.



Figure 3. Glutathione peroxidase (GPx) activity in plasma and liver mitochondria of rats. ^{aa}Significantly different from control group at p < .05. Values are the mean \pm SE (n = 5) ^{bb}Significantly different from CCl₄ group at p < .05. PF, Propofol; AT, (alpha-tocopherol; vitamin E). Duplicate bars from left to right represent plasma and liver mitochondria, respectively.

0.001). Coadministration of Vitamin E with CCl_4 decreased GSH in comparison to propofol (p-value < 0.001); Figure 4.

MTT cocentration

In liver mitochondria: administration of CCl_4 decreased viability of cells in comparison to Vitamin E group (p-value < 0.001). Treatment with propofol increased viability of cells in comparison to CCl_4 group (p-value < 0.001). Coadministration of Vitamin E and propofol with CCl_4 increased viability of mitochondrial cell when compared to CCL_4 respectively(p-value < 0.001, p-value = 0.002); Figure 5.

Discussion

Overall results of the present subchronic

model of toxicity confirmed potential of CCl_4 in induction of oxidative injuries and mitochondrial damage. Results indicated that anti-oxidant enzymes GPx, SOD and oxidative marker such as LPO are stimulated in response to exposure to CCl_4 , but their defense was not sufficient to overcome offending free radicals. As observed in the present results, the GSH is also reduced by CCl_4 . Fascinatingly, propofol indicate antioxidant properties as well as vitamin E were able to attenuate CCl_4 induced changes in almost most tested biomarkers.

In other words, CCl_4 induced oxidative stress by increasing free radicals like induction of cellular damage shown as increase of LPO and reduction of GSH, finally affecting antioxidant enzymes activities. The amelioration effect of propofol may be due to chemically



Figure 4. Reduced glutathione (GSH) in plasma and liver mitochondria of rats. ^{as}Significantly different from control group at p < .05. Values are the mean \pm SE (n = 5) ^{bb}Significantly different from CCl₄ group at p < .05. PF, Propofol; AT, (alpha-tocopherol; vitamin E). Duplicate bars from left to right represent plasma and liver mitochondria, respectively.



Figure 5. Effect of propofol and CCl4 on percent (of control) mitochondrial viabilities of rat liver. ^{aa}Significantly different from control group at p < .05. Values are the mean \pm SE (n = 5) ^{bb}Significantly different from CCl4 group at p < .05. PF, Propofol; AT,(alpha-tocopherol; vitamin E).

similar to endogenous anti-oxidant a-tocopheral (Vitamin E) and theoretically should demonstrate similar properties (20). Numerous studies have demonstrated antioxidant effects of propofol in-vitro (5, 6, 21) and in-vivo (7), but actions of propofol on different cells are varied and multiple mechanisms may be involved. This result indicates reduction of LPO in propofol treatment group induced by CCl₄. The anti-oxidant effects of propofol may also be due to its ability to attenuate the formation of lipid peroxides (22), to induce the expression of anti-oxidant enzyme heme oxygenase-1 (6), to decrease the expression of nitric oxide synthase (NOS), (23) and to stabilize the mitochondrial membrane (24). Our findings showed that, propofol reduced oxidative biomarkers against CCl₄ toxicity in plasma and liver mitochondria. We determined that the propofol, mitochondrial toxicity significantly decreased, whereas it increased significantly in liver mitochondria following CCl₄ administration. We think it is due to anti-oxidant properties of propofol in liver mitochondria. Propofol was also shown to promote mitochondrial function by stabilizing the transmembrane electrical potential (25, 26) and inhibiting mitochondrial permeability transition pore opening (27), both contributing to suppression of mitochondrion-dependent apoptotic signaling (28). These findings

indicate that the toxic stress of CCl_4 and the protective effects of propofol extensively involve mitochondrial viability (Figure 5). The injuries induced by CCl_4 are resulted from free radicals through lipid peroxidation of cell membranes, reduces anti-oxidant enzyme and anti-oxidant substrates to induce oxidative stress that is an important factor in acute and chronic injuries in various tissues (15). The low levels of LPO in the groups receiving propofol in our study suggest that propofol prevents the lipid peroxidation caused by CCl_4 (Figure 1).

Oxidative stress may result in overproduction of oxygen free-radical precursors and/or decreased efficiency of the anti-oxidant system. CCl₄ and oxygen free-radical generation impaired is associated with glutathione metabolism, alterations in the anti-oxidant status (29). However, oxidative stress through generation of reactive oxygen species (ROS) plays an important role in producing liver damage and initiating hepatic fibrogenesis. Oxidative disruption of lipids, proteins, and DNA induces necrosis and apoptosis of hepatocytes, and amplifies the inflammatory response, resulting in initiation of fibrosis. Additionally, ROS stimulate the production of profibrogenic mediators from Kupffer cells and circulating inflammatory cells (30). In present study propofol inhibition of ROS formation, scavenging ROS, or interfering

with ROS pathogenic signaling pathways might be the potential ways to protect mitochondrial cells from dysfunction by CCl₄. While the contribution of Complex III to mitochondrial ROS production in the absence of respiratory inhibitors remains unresolved, two known effects of mitochondrial Ca⁺² could conceivably promote ROS production at this site. One of the key events that cause mitochondrial injury is an abnormal increase in intracellular Ca⁺² (31, 32). It is well known that mtPT can be mediated by a concerted action between Ca²⁺ and ROS, where ROS promote the oxidation and crosslinkage of mitochondrial membrane protein thiol groups such as GSH (33). The mechanisms of Ca⁺²-mediated mitochondrial damage involves activation of degradative enzymes, e.g., calpain proteases and phospholipases, or enzymes that generate reactive oxygen species (ROS) capable of oxidatively modifying mitochondrial proteins and lipids (34, 35). Reduced glutathione is a substrate for mitochondrial matrix glutathione peroxidase, an anti-oxidant enzyme that detoxifies H₂O₂ and other peroxides (36-38). The results of our present investigation showed that 0.2 mL/Kg CCl₄ administration in rats caused significant induction in the activity of anti-oxidant enzymes and decreased GSH. GSH is a critical endogenous anti-oxidant involved in cell defense (39). Although GSH tends to be conserved preferentially in mitochondria, the concentration of GSH in mitochondria may be most relevant for the disposal of ROS and cell fate (40-42). The intracellular distribution of GSH, particulary interms of relative amounts of GSH and disulfide (GSSG), invarious compartments remains unclear. It is a evident, however, that GSH is present in cytosolic, mitochondrial and nuclear pools, can distribute to the nucleous under various condition and is involved in the redox regulation of signal transduction (43-45). Decrease in GSH activity during CCl₄ toxicity might be due to the decreased availability of GSH resulted during the enhanced oxidative damage (46-48). Improvement of blood and liver mitochondrial GSH levels in rats treated with propofol in comparison to CCl₄ administration further demonstrated the anti-oxidant properties of this drug (Figure 4). This effect is established via activating the glutathione anti-oxidant system (49, 50).

ROS are produced in normal metabolic processes and are inactivated by endogenous antioxidant systems (51). The primary ROS produced by mitochondria is superoxide [89]. This highly reactive free radical is extremely short-lived (52-54) and dismutates either spontaneously or with the help of the mitochondrial superoxide dismutase forming the more ROS, H_2O_2 (55, 56). According to our opinion, the significant decrease of GPx and SOD activity in Groups Vitamin when compared to CCl₄ group may be explained with the suppression of physiologic ROS production by anti-oxidant agents (Figure 2, 3).

Recently in-vitro studies showed that propofol could effectively suppress apoptotic signaling and prevent apoptotic death of myocardial cells encountering fatal stimuli (6, 57, 58). In experimental hearts, propofol inhibited mitochondrial permeability transition (27) and ameliorated ischemia-reperfusion injury (59-61). In fact, sustained hepatic inflammation provoked by long-term treatment with CCl₄ is believed to be through NF-KB pathway (62, 63). NF- κ B is a transcription factor consisting of p65 and p50 subunits of the Rel protein family that regulates host inflammatory and immune responses by increasing the expression of specific cellular genes (64, 65). Released activated NF- κB complex tranlocates to the nucleus, where it regulates gene expression by binding to kB binding sites (61, 66).

These findings support the premise that propofol can guard against the sequences of oxidative stress. Hence, the anti-oxidant properties of propofol are involved in its hepatoprotective mechanism. The powerful antioxidant activity of propofol has been examined previously in different oxidative stress situations (67). Interestingly, as evidenced by MTT assay, the present data confirm that CCl_4 -induced mitochondrial dysfunction can be improved by propofol. However, the exact molecular mechanisms of a putative protective role of propofol in hepatoprotection should be further investigated.

Acknowledgement

This study was supported by a grant from

proposal numbered 90007921 at Iran National Science Foundation (INSF).

References

- Marik PE. Propofol: an immunomodulating agent. *Pharmacotherapy. J. Hum. Pharm. Drug Ther.* (2005) 25: 28-33.
- (2) Tramer M, Moore A and McQuay H. Propofol anaesthesia and postoperative nausea and vomiting: quantitative systematic review of randomized controlled studies. *Br. J. Anaes*. (1997) 78: 247-255.
- (3) Schaer H. Propofol infusion for the maintenance of short-term anesthesia. *Anaes*. (1988) 37: 187-192.
- (4) Ansley DM, Lee J, Godin DV, Garnett ME and Qayumi AK. Propofol enhances red cell antioxidant capacity in swine and humans. *Can. J. Anest.* (1998) 45: 233-239.
- (5) Tesauro M, Thompson W and Moss J. Effect of staurosporine-induced apoptosis on endothelial nitric oxide synthase in transfected COS-7 cells and primary endothelial cells. *Cell Death Differen*. (2005) 13: 597-606.
- (6) Xu J and Wang Y. Propofol attenuation of hydrogen peroxide-mediated oxidative stress and apoptosis in cultured cardiomyocytes involves haeme oxygenase-1. *Eur. J. Anaes.* (2008) 25: 395-402.
- (7) Kobayashi K, Yoshino F, Takahashi SS, Todoki K, Maehata Y, Komatsu T, Yoshida K and Lee MC Direct assessments of the anti-oxidant effects of propofol medium chain triglyceride/long chain triglyceride on the brain of stroke-prone spontaneously hypertensive rats using electron spin resonance spectroscopy. *Anest.* (2008) 109: 426-435.
- (8) Chen J, Gu Y, Shao Z, Luo J and Tan Z. Propofol protects against hydrogen peroxide-induced oxidative stress and cell dysfunction in human umbilical vein endothelial cells. *Mole. Cell Biochem.* (2010) 339: 43-54.
- (9) Wang B, Shravah J, Luo H, Raedschelders K, Chen DDY and Ansley DM. Propofol protects against hydrogen peroxide-induced injury in cardiac H9c2 cells via Akt activation and Bcl-2 up-regulation. *Biochem. Biophys. Res. Commun.* (2009) 389: 105-111.
- (10) Luo T and Xia Z. A small dose of hydrogen peroxide enhances tumor necrosis factor-alpha toxicity in inducing human vascular endothelial cell apoptosis: reversal with propofol. *Anest. Analg.* (2006) 103: 110-116.
- (11) Xia Z, Luo T, Liu H, Wang F, Irwin MG and Vanhoutte PM. L-arginine enhances nitrative stress and exacerbates tumor necrosis factor-[alpha] toxicity to human endothelial cells in culture: prevention by propofol. J. Cardiol. Pharm. (2010) 55: 358.
- (12) Abdollahi M, Ranjbar A, Shadnia S, Nikfar S and Rezaie A. Pesticides and oxidative stress: a review. *Med. Sci. Monitor* (2004) 10: 141.
- (13) Ranjbar A, Ghahremani MH, Sharifzadeh M, Golestani A, Ghazi-Khansari M, Baeeri M and Abdollahi M. Protection by pentoxifylline of malathion-induced toxic stress and mitochondrial damage in rat brain. *Hum. Experi. Toxicol.* (2010) 29: 851-864.
- (14) Soleimani H, Ranjbar A, Baeeri M, Mohammadirad

A, Khorasani R, Yasa N and Abdollahi M. Rat plasma oxidation status after *Nigella Sativa* L. botanical treatment in CCl₄-treated rats. *Toxicol. Mech. Meth.* (2008) 18: 725-731.

- (15) Szymonik-Lesiuk S, Czechowska G, Stryjecka-Zimmer M, Słomka M, Madro A, Celiński K and Wielosz M. Catalase, superoxide dismutase, and glutathione peroxidase activities in various rat tissues after carbon tetrachloride intoxication. J. Hep. Bili. Pan. Sur. (2003) 10: 309-315.
- (16) Ghazi-Khansari M, Mohammadi-Bardbori A and Hosseini MJ. Using janus green B to study paraquat toxicity in rat liver mitochondria. *Ann. New York Acad. Sci.* (2007) 1090: 98-107.
- (17) Moore K and Roberts LJ. Measurement of lipid peroxidation. *Free Rad. Res.* (1998) 28: 659-671.
- (18) Kruger NJ. The Bradford method for protein quantitation. *Methods Mol. Biol.* (1994) 32: 9-15.
- (19) Morgan DML. Tetrazolium (MTT) assay for cellular viability and activity. *Method Molec. Biol. Clif. Totow.* (1997) 79: 179-183.
- (20) Bayona NA, Gelb AW, Jiang Z, Wilson JX, Urquhart BL and Cechetto DF. Propofol neuroprotection in cerebral ischemia and its effects on low-molecularweight antioxidants and skilled motor tasks. *Anest.* (2004) 100: 1151.
- (21) Wang B, Luo T, Chen D and Ansley DM. Propofol reduces apoptosis and up-regulates endothelial nitric oxide synthase protein expression in hydrogen peroxide-stimulated human umbilical vein endothelial cells. *Anest. Analg.* (2007) 105: 1027-1033.
- (22) Kahraman S, Kilinc K, Dal D and Erdem K. Propofol attenuates formation of lipid peroxides in tourniquetinduced ischaemia-reperfusion injury. *Brit. J. Anaes.* (1997) 78: 279-281.
- (23) Goode HF, Cowley HC, Walker BE, Howdle PD and Webster NR. Decreased antioxidant status and increased lipid peroxidation in patients with septic shock and secondary organ dysfunction. *Crit. Care Med.* (1995) 23: 646-651.
- (24) Eriksson O. Effects of the general anaesthetic Propofol on the Ca²⁺-induced permeabilization of rat liver mitochondria. *FEBS Lett.* (1991) 279: 45-48.
- (25) Branca D, Roberti MS, Lorenzin P, Vincenti E and Scutari G. Influence of the anesthetic 2, 6-diisopropylphenol on the oxidative phosphorylation of isolated rat liver mitochondria. *Biochem. Pharm.* (1991) 42: 87-90.
- (26) Tsujimoto Y and Shimizu S. Role of the mitochondrial membrane permeability transition in cell death. *Apop.* (2007) 12: 835-840.
- (27) Javadov SA, Lim KHH, Kerr PM, Suleiman MS, Angelini GD and Halestrap AP. Protection of hearts from reperfusion injury by propofol is associated with inhibition of the mitochondrial permeability transition. *Card. Res* (2000) 45: 360-369.
- (28) Crompton M. The mitochondrial permeability transition pore and its role in cell death. *Biochem. J.* (1999) 341: 233.

- (29) Gilbert MT, Haselkorn T, Bunce M, Sanchez JJ, Lucas SB, Jewell LD, Van Marck E and Worobey M. The isolation of nucleic acids from fixed, paraffinembedded tissues–which methods are useful when? *PLoS One.* (2007) 2: 537.
- (30) Friedman SL. Mechanisms of hepatic fibrogenesis. *Gastro*. (2008) 134: 1655-1669.
- (31) Schild L, Huppelsberg J, Kahlert S, Keilhoff G and Reiser G. Brain mitochondria are primed by moderate Ca²⁺ rise upon hypoxia/reoxygenation for functional breakdown and morphological disintegration. *J. Biol. Chem.* (2003) 278: 25454-25460.
- (32) Abo-Hashema KAH, Cake MH and Potter IC. Liver mitochondria, confirmed as intact by complete suppression of succinate uptake and oxidation, possess a carnitine palmitoyltransferase I that is totally inhibited by malonyl CoA. *Biochem. Biophys. Res. Commun.* (1999) 258: 778-783.
- (33) Maciel EN, Vercesi AE and Castilho RF. Oxidative stress in Ca²⁺-induced membrane permeability transition in brain mitochondria. J. Neurochem. (2002) 79: 1237-1245.
- (34) Starkov AA, Chinopoulos C and Fiskum G. Mitochondrial calcium and oxidative stress as mediators of ischemic brain injury. *Cell Cal.* (2004) 36: 257-264.
- (35) Borutaite V. Mitochondria as decision-makers in cell death. *Environm. Mol. Mut.* (2010) 51: 406-416.
- (36) Andreyev AY, Kushnareva YE and Starkov A. Mitochondrial metabolism of reactive oxygen species. *Biochem.* (2005) 70: 200-214.
- (37) Marí M, Morales A, Colell A, García-Ruiz C and Fernández-Checa JC. Mitochondrial glutathione, a key survival antioxidant. *Antioxid. Redox. Signal.* (2009) 11: 2685-2700.
- (38) Shlafer M, Myers CL and Adkins S. Mitochondrial hydrogen peroxide generation and activities of glutathione peroxidase and superoxide dismutase following global ischemia. J. Mol. Cell Card. (1987) 19: 1195-206.
- (39) Fernández-Checa J, García-Ruiz C, Colell A, Morales A, Marí M, Miranda M and Ardite E. Oxidative stress: role of mitochondria and protection by glutathione. *Biof.* (2008) 8: 7-11.
- (40) Griffith OW and Meister A. Origin and turnover of mitochondrial glutathione. *Proceed. Nat. Acad. Sci.* USA. (1985) 82: 4668.
- (41) Kurosawa K, Hayashi N, Sato N, Kamada T and Tagawa K. Transport of glutathione across the mitochondrial membranes. *Biochem. Biophy. Res. Commun.* (1990) 167: 367-372.
- (42) von Montfort C, Matias N, Fernandez A, Fucho R, Conde de la Rosa L, Martinez-Chantar ML, Mato JM, Machida K, Tsukamoto H, Murphy MP, Mansouri A, Kaplowitz N, Garcia-Ruiz C and Fernandez-Checa JC. Mitochondrial GSH determines the toxic or therapeutic potential of superoxide scavenging in steatohepatitis. J. Hep. (2012) 57: 852-859.
- (43) Voehringer D, McConkey D, McDonnell T, Brisbay

S and Meyn R. Bcl-2 expression causes redistribution of glutathione to the nucleus. *Proceed. Nat. Aca. Sci.* (1998) 95: 2956-2960.

- (44) Briviba K, Fraser G, Sies H and Ketterer B. Distribution of the monochlorobimane-glutathione conjugate between nucleus and cytosol in isolated hepatocytes. *Biochem. J.* (1993) 294: 631.
- (45) Nunes C, Barbosa RM, Almeida L and Laranjinha J. Nitric oxide and DOPAC-induced cell death: from GSH depletion to mitochondrial energy crisis. *Mol. Cell. Neurosci.* (2011) 48: 94-103.
- (46) Khan RA, Khan MR, Ahmed M, Sahreen S, Shah NA, Shah MS, Bokhari J, Rashid U, Ahmad B and Jan S. Hepatoprotection with a chloroform extract of *Launaea procumbens* against CCl₄-induced injuries in rats. *BMC Complem. Alter. Med.* (2012) 12: 114.
- (47) Mujeeb M, Khan SA, Aeri V and Ali B. Hepatoprotective Activity of the Ethanolic Extract of Ficus carica Linn. Leaves in Carbon Tetrachloride-Induced Hepatotoxicity in Rats. *Iran. J. Pharm. Sci.* (2011) 10: 301-306.
- (48) Balbis E, Patriarca S, Furfaro AL, Millanta S, Sukkar SG, Marinari UM, Pronzato MA, Cottalasso D and Traverso N. Whey proteins influence hepatic glutathione after CCl₄ intoxication. *Toxicol. Indus. Health.* (2009) 25: 325-328.
- (49) De La Cruz J, Zanca A, Carmona J and de la Cuesta FS. The effect of propofol on oxidative stress in platelets from surgical patients. *Anest. Analg.* (1999) 89: 1050.
- (50) Oztürk E, Demirbilek S, Köroğlu A, But A, Begeç ZO, Gülec M, Akyol O and Ersoy MO. Propofol and erythropoietin antioxidant properties in rat brain injured tissue. *Prog. Neuro-Psychopharm. Biol. Psych.* (2008) 32: 81-86.
- (51) Halliwell B. Free radicals and antioxidants: a personal view. Nut. Rev. (1994) 52: 253-565.
- (52) Dionisi O, Galeotti T, Terranova T and Azzi A. Superoxide radicals and hydrogen peroxide formation in mitochondria from normal and neoplastic tissues. *Biochim. Biophys. Acta*. (1975) 403: 292.
- (53) Buettner GR. Superoxide dismutase in redox biology: the roles of superoxide and hydrogen peroxide. *Antican. Age Med. Chem.* (2011) 11: 341.
- (54) Zhang L, Yu L and Yu CA. Generation of Superoxide Anion by Succinate-Cytochromec Reductase from Bovine Heart Mitochondria. J. Biol. Chem. (1998) 273: 33972-33976.
- (55) Murphy MP. How mitochondria produce reactive oxygen species. *Biochem. J.* (2009) 417: 1.
- (56) Sawyer DT and Valentine JS. How super is superoxide? Acco. Chem. Res. (1981) 14: 393-400.
- (57) Kim HS, Chang WC, Hwang KC, Choi IG and Park WK. Effect of propofol on calcium homeostasis in hypoxia-reoxygenated neonatal rat cardiomyocytes. *Eur. J. Pharm.* (2008) 594: 139-145.
- (58) Noh HS, Shin IW, Ha JH, Hah YS, Baek SM and Kim DR. Propofol protects the autophagic cell death induced by the ischemia/reperfusion injury in rats. *Mol. Cells.* (2010) 30: 455-460.

- (59) Dehkordi ME, Razavi SS and Momenzadeh S. A Comparison between Sedative Effect of Propofol-Fentanyl and Propofol-Midazolam Combinations in Microlaryngeal Surgeries. *Iran. J. Pharm. Sci.* (2012) 11: 287-94.
- (60) Xia W, Liu Y, Zhou Q, Tang Q and Zou H. Comparison of the effects of propofol and midazolam on inflammation and oxidase stress in children with congenital heart disease undergoing cardiac surgery. *Yon. Med. J.* (2011) 52: 326-332.
- (61) Lai HC, Yeh YC, Wang LC, Ting CT, Lee WL, Lee HW, Wang KY, Wu A, Su CS and Liu TJ. Propofol ameliorates doxorubicin-induced oxidative stress and cellular apoptosis in rat cardiomyocytes. *Toxicol. App. Pharm.* (2011) 257: 437-448.
- (62) Orfila C, Lepert JC, Alric L, Carrera G, Béraud M and Pipy B. Immunohistochemical distribution of activated nuclear factor κB and peroxisome proliferatoractivated receptors in carbon tetrachloride-induced chronic liver injury in rats. *Histochem. Cell Bio*. (2005) 123: 585-593.
- (63) Son G, Iimuro Y, Seki E, Hirano T, Kaneda Y and

Fujimoto J. Selective inactivation of NF- κ B in the liver using NF- κ B decoy suppresses CCl₄-induced liver injury and fibrosis. *Am. J. Physiol. Gastro. Liver Physiol.* (2007) 293: 631-639.

- (64) Ali S and Mann DA. Signal transduction via the NF-κB pathway: a targeted treatment modality for infection, inflammation and repair. *Cell Biochem. Func.* (2004) 22: 67-79.
- (65) Hamdy N and El-Demerdash E. New therapeutic aspect for carvedilol: Antifibrotic effects of carvedilol in chronic carbon tetrachloride-induced liver damage. *Toxicol. Appl. Pharm.* (2012) 261: 292-299.
- (66) Hayden M, West A and Ghosh S. NF-κB and the immune response. *Oncogene* (2006) 25: 6758-6780.
- (67) Kostopanagiotou GG, Grypioti AD, Matsota P, Mykoniatis MG, Demopoulos CA, Papadopoulou-Daifoti Z and Pandazi A. Acetaminophen-induced liver injury and oxidative stress: protective effect of propofol. *Eur. J. Anaest.* (2009) 26: 548-553.

This article is available online at http://www.ijpr.ir