

The Effect of Chronic Administration of Safranal on Systolic Blood Pressure in Rats

Mohsen Imenshahidi^a, Bibi Marjan Razavi^b, Ayyoob Faal^c, Ali Gholampoor^c,
Seyed Mehran Mousavi^c and Hossein Hosseinzadeh^{a*}

^aPharmaceutical Research Center, Department of Pharmacodynamics and Toxicology, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran. ^bTargeted Drug Delivery Research Center, Department of Pharmacodynamics and Toxicology, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran. ^cSchool of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran.

Abstract

Safranal, the main component of *Crocus sativus* essential oil, exhibits different pharmacological activities. In this study, the effects of safranal, on blood pressure of normotensive and desoxycorticosterone acetate (DOCA) - salt induced hypertensive rats in chronic administration were investigated. Three doses of safranal (1, 2 and 4 mg/Kg/day) and spironolactone (50 mg/Kg/day) were administrated to the different groups of normotensive and hypertensive rats (at the end of 4 weeks treatment by DOCA-salt) for Five weeks. Then the effects of safranal on mean systolic blood pressure (MSBP) and heart rate (HR) were evaluated using tail cuff method. The duration of effect of safranal on SBP, was also evaluated. Our results indicated that chronic administration of safranal could reduce the MSBP in DOCA salt treated rats in a dose dependent manner. Safranal did not decrease the MSBP in normotensive rats. The data also showed that antihypertensive effects of safranal did not persist. In summary, our results showed that safranal exhibits antihypertensive and normalizing effect on BP in chronic administration.

Keywords: *Crocus sativus*; Saffron; Safranal; Systolic blood pressure; Tail cuff.

Introduction

Crocus sativus L., a member of Iridaceae family, is commonly known as saffron. It is widely cultivated in Iran and other countries (1). Safranal, the main ingredient of *C. sativus* essential oil, is thought to be responsible for the saffron unique odor (2). In addition, safranal is also known as a potent antioxidant (3). It was reported that safranal has anticonvulsant (4), antidepressant (5), antianxiety and hypnotic

(6), antitussive (7), antinociceptive (8) and anti-ischemia (9) effects and could improve memory and learning ability in rats (10). The protective effect of safranal on diazinon induced oxidative stress has been shown recently (11). The effect of saffron and its components on cardiovascular system have also been established in some studies. It was indicated that aqueous-ethanol extract of *C. sativus*, exhibits a potent inhibitory effect on heart rate and contractility of guinea pig heart via calcium channel-blocking effect (12). Moreover, the hypotensive effect of *C. sativus* petals extract in rats has been shown previously

* Corresponding author:

E-mail: hosseinzadehh@mums.ac.ir

Table 1. Summary of selected groups.

Groups	DOCA	Normal saline	Safranal	Spironolactone
1		*		
2	*	*		
3,4 and 5	*		*	
6	*			*
7		*	*	

(13). The results of our previous study revealed that the aqueous extract of saffron stigma as well as two major constituents of this plant, crocin and safranal, has hypotensive properties in normotensive and hypertensive anaesthetized rats (14). Although the effect of this plant in lowering blood pressure have been shown previously, but there has not been any study about the effect of safranal on blood pressure in chronic administration. Thus, in this study the effects of chronic administration of safranal, on blood pressure of normotensive and desoxycorticosterone acetate (DOCA)-salt induced hypertensive rats were investigated.

Experimental

Animal and chemicals

Adult male Wistar rats (weight 250–300 g) were provided by animal center (School of Pharmacy, Mashhad University of Medical Sciences). They were maintained on a 12 h light/dark cycle and at a temperature of 23 ± 1 °C with free access to food and water. These conditions were maintained constant throughout the experiments. The experiments were performed under the Animals (scientific procedures) Act of 1986 and conform to the National Institutes of Health guidelines for the use of experimental animals. Safranal and Tween-80 were obtained from Merck (Germany) and BDH Chemicals (Heidelberg, Germany). 5% Tween-80 was used as negative control.

Induction of experimental hypertension

Desoxycorticosterone acetate (DOCA)-salt (20 mg/Kg, twice weekly, for 4 weeks, s.c.) and NaCl (1%) in rat's drinking water were used for induction of hypertension (14). Rats were randomly divided into 7 groups. 1) Saline injected (0.5 mL/Kg, twice weekly, s.c.,

for 4 weeks), this treatment was continued for another five weeks, 2) (DOCA)-salt (20 mg/kg, twice weekly, for 4 weeks, s.c.), DOCA treatment was continued by *i.p.* injection of 0.5 mL/Kg normal saline for another five weeks, 3, 4 and 5) (DOCA)-salt (20 mg/Kg, twice weekly, for 4 weeks, s.c.), DOCA treatment was continued by *i.p.* injection of 1, 2 and 4 mg/Kg/day safranal for another five weeks, after that safranal injection was stopped but DOCA injection was continued for another two weeks, 6) (DOCA)-salt (20 mg/Kg, twice weekly, for 4 weeks, s.c.), DOCA treatment was continued by *i.p.* injection of 50 mg/Kg/day spironolactone for another five weeks, after that spironolactone injection was stopped but DOCA injection was continued for another two weeks, 7) Saline injected (0.5 mL/Kg, twice weekly, s.c., for 4 weeks), saline treatment was continued by *i.p.* injection of 4 mg/Kg safranal for another five weeks. All groups consisted of six rats. Table 1 describes the different groups that were selected for this study.

Hypotensive activity

Four, nine and eleven weeks after the first saline or DOCA treatment, SBP was measured using tail cuff method in all groups as described by Lorenz (15). Briefly, three days before the last treatment, the training of rats in different groups for indirect SBP measurements was started. This training consisted of the regular handling of the animals and getting used to the restraining cage and the tail-cuff. Rats were heated for approximately 15 minutes at 30-32 °C to increase blood flow to the tail. After that, animals were placed in small restraining cages with a cuff around the end of proximal of the tail. After placing of the cuff, a pulse transducer was used around the end of the tail. Then the




Figure 1. Hypertension induced by Desoxycorticosterone acetate (DOCA)-salt after 4 weeks. Each value is the mean \pm SEM of six experiments, *** $P < 0.001$ vs normal saline treated rats. One way ANOVA, Tukey Krumer test.

tail cuff was inflated using the related button on the NIBP (Non-Invasive Blood Pressure) controller apparatus and acquisition data were performed by a computerized system Power Lab (ADInstruments, v 5.4.2). The mean values of five BPs and HRs readings were used for each animal.

Statistical analysis

Results are expressed as mean \pm SEM. Statistical analysis was performed with ANOVA followed by Tukey–Kramer test to compare the differences between means. Differences were considered statistically significant when $P < 0.05$.

Results

Effect of DOCA on SBP

In DOCA treated rats, MSBP significantly increased in comparison with normal saline treated (normotensive) rats ($P < 0.001$) (Figure 1).

Effects of safranal on normotensive and hypertensive rats after nine weeks

As shown in Figure 2 the injection of safranal (1, 2 and 4 mg/Kg) decreased the MSBP in hypertensive animals ($P < 0.001$, respectively), but in normotensive rats, safranal did not reduce the MSBP. The hypotensive effect of safranal was similar to that of spironolactone in different doses.




Figure 2. Mean systolic blood pressure (MSBP) in response to various doses of safranal in normotensive and hypertensive rats at the end of nine weeks. Each value is the mean \pm SEM of six experiments. One-way ANOVA, Tukey Krumer, *** $P < 0.001$ vs DOCA plus normal saline treated rats.




Figure 3. Evaluation of the duration hypotensive effect of safranal. Each value is the mean \pm SEM of six experiments. One-way ANOVA, Tukey Krumer, **P<0.001 vs DOCA plus normal saline treated rats.

Evaluation of duration effect of safranal on SBP

As shown in Figure 3, although the hypotensive effect of safranal did not persist (97.00 ± 5.00 and 127.00 ± 7.00 , before and after stopping the safranal injection, respectively), but the decrease in SBP induced by safranal, did not reach to that of DOCA treated rats after stopping the injection (P<0.01).

Effect of safranal on heart rate

Safranal caused a reduction in heart rate, but the effect was not significant (Figure 4).

Discussion

Deoxycorticosterone acetate (DOCA)-salt is an agent commonly used to induce hypertension in experimental animals (14). Our results showed that DOCA-salt significantly




Figure 4. Effect of safranal on heart rate in hypertensive rats (before and after injection). Each value is the mean \pm SEM of six experiments. T student test and one- way ANOVA.

induced hypertension in comparison with saline group at the end of 4 weeks treatment. Chronic administration of safranal reduced the increase of MSBP induced by DOCA, but this hypotensive effect was not observed in normotensive rats.

Previous studies revealed that saffron and its constituents exhibit vasodilatory effects. For examples, a potent relaxant effect of safranal on smooth muscles of guinea pigs has been shown (16), so that it might be concluded that hypotensive effect of safranal in chronic treatment is attributed to the inhibitory effect on smooth muscles through blocking of calcium channel or inhibition of sarcoplasmic reticulum Ca^{2+} release into cytosol. Also, it was reported that aqueous and ethanolic extracts of saffron petals, reduced the MABP in anaesthetized rats (13). Moreover it was indicated that intravenous injection of aqueous extract of saffron stigma and two major constituents of this plant exhibit hypotensive effects in normotensive and hypertensive anaesthetized rats and the effect of safranal on lowering blood pressure was more than crocin, so it could be suggested that safranal is more important than crocin in hypotensive properties of saffron (14). According to LD50 values, safranal was low-toxic in acute intraperitoneal route and practically non-toxic in acute oral administration in both mice and rats (17). Our results showed safranal did not cause reflex tachycardia, so it might be concluded that both heart function and blood vessels contractility are affected by safranal (14).

Based on the previous study, one of the mechanisms involved in anti hypertensive effect of safranal like *Asperugo procumbens* (18) might be related to its effect on GABA (A)-benzodiazepine receptor complex (17). Because benzodiazepines, in pre-anesthetic doses, reduce blood pressure through decreasing peripheral resistance or cardiac output (18).

It is indicated that administration of DOCA-salt causes pathophysiological and biochemical changes so that DOCA-salt hypertensive rats, could provide an animal model of oxidative and inflammatory stress in the cardiovascular system (20). Therefore, the DOCA-salt experiment can provide an appropriate model to evaluate of anti-oxidative or anti-inflammatory responses of natural or synthetic compounds on cardiovascular system. This also provides opportunities for the

development of novel therapeutic agents for management of chronic cardiovascular disease (21). Therefore, it could be concluded that the antihypertensive effects of safranal could be related partly due to their antioxidant properties (3). It is well known that DOCA induced hypertension causes an endothelial dysfunction in the isolated aortic rings as well as in the perfused mesenteric bed (22). Since safranal decreased SBP in hypertensive rats, our results probably show that the vasodilatory effect of safranal was endothelium dependent.

Spiroolactone, known as potassium-sparing diuretics, inhibits the effects of aldosterone by competing for intracellular mineralocorticoid receptors in the cortical collecting duct. This decreases the reabsorption of sodium and water, and decreasing the secretion of potassium (23). In this study, spiroolactone was used as a positive control. Our results showed that the antihypertensive effect of safranal at the highest dose was as much as spiroolactone at the end of nine weeks. It is likely that the hypotensive effect of safranal may be due to the saffron diuretic effect (1).

To evaluate the duration of effects of safranal on reducing SBP, the safranal injection was stopped at the end of nine weeks but DOCA injections were continued for another two weeks. The data also revealed that although the hypotensive effect of the highest dose of safranal did not persist completely and SBP increased again after stopping the injection, but SBP did not restored to that of DOCA treated rats, this might be related to the incomplete elimination of safranal.

Conclusion

In summary, our results indicated that chronic administration of safranal could reduce the MSBP in DOCA salt treated rats. Therefore, safranal exhibits antihypertensive and normalizing effect on BP.

Acknowledgments

The authors are thankful to the Vice Chancellor of Research, Mashhad University of Medical Sciences for financial support.

References

- (1) Ríos J, Recio M, Giner R and Manez S. An update review of saffron and its active constituents. *Phytother. Res.* (1996) 10: 189-193.
- (2) Rezaee R and Hosseinzadeh H. Safranal: From an aromatic natural product to a rewarding pharmacological agent. *Iran. J. Basic. Med. Sci.* (2013) 16: 12-26.
- (3) Assimopoulou AN, Sinakos Z and Papageorgiou VP. Radical scavenging activity of *Crocus sativus* L extract and its bioactive constituents. *Phytother. Res.* (2005) 19: 997-1000.
- (4) Hosseinzadeh H and Talebzadeh F. Anticonvulsant evaluation of safranal and crocin from *Crocus sativus* in mice. *Fitoterapia* (2005) 76: 722-724.
- (5) Hosseinzadeh H, Karimi G and Niapoor M. Antidepressant effects of *Crocus sativus* stigma extracts and its constituents, crocin and safranal, in mice. *J. Med. Plants* (2004) 3: 48-58.
- (6) Hosseinzadeh H and Noraei NB. Anxiolytic and hypnotic effect of *Crocus sativus* aqueous extract and its constituents, crocin and safranal, in mice. *Phytother. Res.* (2009) 23: 768-774.
- (7) Nemati H, Boskabady MH and Ahmadzadeh Vostakolaei H. Stimulatory effect of *Crocus sativus* (saffron) on β_2 -adrenoceptors of guineapigtrachealchains. *Phytomed.* (2008) 15: 1038-1045.
- (8) Amin B and Hosseinzadeh H. Evaluation of aqueous and ethanolic extracts of saffron, *Crocus sativus* L and its constituents, safranal and crocin in allodynia and hyperalgesia induced by chronic constriction injury model of neuropathic pain in rats. *Fitoterapia* (2012) 83: 888-895.
- (9) Hosseinzadeh H and Sadeghnia HR. Safranal, a constituent of *Crocus sativus* (saffron), attenuated cerebral ischemia induced oxidative damage in rat hippocampus. *J. Pharm. Pharm. Sci.* (2005) 8: 394-399.
- (10) Hosseinzadeh H and Ziaei T. Effects of *Crocus sativus* stigma extract and its constituents, crocin and safranal, on intact memory and scopolamine-induced learning deficits in rats performing the Morris water maze task. *J. Med. Plants* (2006) 5: 40-50.
- (11) Hariri A, Moallem S, Mahmoudi M and Hosseinzadeh H. The effect of crocin and safranal, constituents of saffron, against subacute effect of diazinon on hematological and genotoxicity indices in rats. *Phytomed.* (2011) 18: 499-504.
- (12) Boskabady MH, Shafei MN, Shakiba A and Sang Sefidi H. Effect of aqueous-ethanol extract from *Crocus sativus* (saffron) on guinea-pig isolated heart. *Phytother. Res.* (2008) 22: 330-334.
- (13) Fatehi M, Rashidabady T and Hassanabad ZF. Effects of petals extracts of saffron on rat blood pressure and on responses induced by electrical field stimulation in the rat isolated vas deferens and guinea-pigileum. *J. Ethnopharmacol.* (2003) 84: 199-203.
- (14) Imenshahidi M, Hosseinzadeh H and Javadpour Y. Hypotensive effect of aqueous saffron extract (*Crocus sativus* L.) and its constituents, safranal and crocin, in normotensive and hypertensive rats. *Phytother. Res.* (2010) 24: 990-994.
- (15) Lorenz JN. A practical guide to evaluating cardiovascular, renal, and pulmonary function in mice. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* (2002) 282: 1565-1582.
- (16) Boskabadi IMH and Aslani MR. Relaxant effect of *crocus sativus* (saffron) on guinea pig tracheal chains and its possible mechanisms. *J. Pharm. Pharmacol.* (2006) 58: 1385-1390.
- (17) Hosseinzadeh H, Shakib SS, Sameni AK and Taghiabadi E. Acute and subacute toxicity of safranal, a constituent of saffron, in mice and rats. *Iran. J. Pharm. Res.* (2013) 12: 93-99.
- (18) Mirshafa SA, Azadbakht M and Ahangar N. Study of antidepressant and sedative-hypnotic activity of hydroalcoholic extract of *Asperugo procumbens* L. aerial parts in mice. *Iran. J. Pharm. Res.* (2013) 12: 529-535.
- (19) Hosseinzadeh H and Sadeghnia H. Protective effect of safranal on pentylenetetrazol-induced seizures in the rat: involvement of GABAergic and opioids systems. *Phytomed.* (2007) 14: 256-262.
- (20) Nugent M, Artru A and Michenfelder J. Cerebral metabolic vascular and protective effects of midazolam maleate: Comparison to diazepam. *Anesthesiol.* (1982) 56: 172-176.
- (21) Lyer A, Chan V and Brown L. The DOCA-Salt hypertensive rat as a model of cardiovascular oxidative and inflammatory stress. *Current. Cardiol. Rev.* (2010) 6: 291-297.
- (22) Fatehi-Hassanabad Z, Fatehi M and Imen Shahidi M. Endothelial dysfunction in aortic rings and mesenteric beds isolated from deoxycorticosterone acetate hypertensive rats: possible involvement of protein kinase C. *Eur. J. Pharmacol.* (2004) 494: 199-204.
- (23) Cheng S, Suzuki K, Sadee W and Harding B. Effects of spironolactone, canrenone and canrenoate-K on cytochrome P450, and 11 β - and 18-hydroxylation in bovine and human adrenal cortical mitochondria. *Endocrinol.* (1974) 99: 1097-1106.

This article is available online at <http://www.ijpr.ir>