

Original Article

The Contents of Sesamol in Iranian Sesame Seeds

Naficeh Sadeghi^a, Mohammad Reza Oveisi^a, Mannan Hajimahmoodi^a,
Behrooz Jannat^{b*}, Masoomeh Mazaheri^a and Sadollah Mansouri^c

^aDrug and Food Control Department, Faculty of Pharmacy, Medical Sciences/University of Tehran, Tehran, Iran. ^bFood and Drug Laboratory Research Center, Ministry of Health and Medical Education, Tehran, Iran. ^cSeed and Plant Improvement Institute, Karaj, Iran.

Abstract

Sesamol is a sesame lignan. Sesame lignans have multiple functions, including antioxidant activity and also seem to have potential as a source of phytoestrogens. This study was conducted to evaluate sesamol contents of 7 brands of Iranian sesame seeds (*Sesamum indicum* L.). The brands were named *Karaj 29*, *Darab 14*, *Ultan*, *Dezful*, *Varamin*, *Branching Naz*, and *Nonbranching Naz*. After method validation, the methanolic extracts of seeds were investigated by HPLC. Their mean of total sesamol was found to be 4.67 ± 0.92 mg/g (ranging between 2.75 and 6.13 mg/g). The brands *Karaj 29* with 5.84 ± 0.25 mg/g, *Dezful* with 5.48 ± 0.08 mg/g, and *Varamin* with 5.4 ± 0.1 mg/g had the highest content, and *Darab 14* with the content of 3.30 ± 0.57 mg/g had the lowest ($p < 0.05$). Iranian sesame can be considered to be a good source of natural antioxidants for medicinal and commercial uses.

Keywords: Sesamol; Lignan; Antioxidant activity; Sesame seed; Analyze; HPLC.

Introduction

Sesame (*Sesamum indicum* L.) seed is one of the most important oil seed crops in the world (1) and is also known as sesamum, gingelly, beniseed, sim-sim, and till. It has been cultivated for centuries, particularly in Asia and Africa, for its high edible oil and protein content (2). It is also considered to be a beneficial food to health (3).

Sesame oil compounds have multiple physiological functions, such as estrogenic activity (4), providing anti-inflammatory functions (5), decreasing blood lipids (6) and arachidonic acid levels (7), and increasing antioxidative ability and γ -tocopherol bioavailability (8).

Recently the development of novel lipid-containing processed foods has increased rancidity caused by lipid autoxidation, posing a serious problem for the keeping quality of such systems (9). However, the oxidative stability of sesame oil is superior to that of other vegetable oils even though it contains nearly 85% unsaturated fatty acids (10). Sesame oil is especially stable because of the presence of unusual compounds known as lignans, comprised of sesamin, sesamolin, sesamol, and γ -tocopherol (11). Some chemicals that occur naturally in plants have begun to receive much attention as safe antioxidants, as they have been consumed by people and animals for years (12).

The present study was undertaken to determine the levels of sesamol lignan in some Iranian sesame seed cultivars and to find the richest one containing this chemical antioxidant.

* Corresponding author:

E-mail: Janatbhr@sina.tums.ac.ir

Experimental

Reagents

The sample of seven different brands of sesame seeds (*Sesamum indicum* L.) were donated by Seed and Plant Improvement Institute, Karaj, Iran.

The brands were named *Karaj 29*, *Darab 14*, *Ultan*, *Dezful*, *Varamin*, *Branching Naz* and *Nonbranching Naz*.

The standard of sesamol was obtained from Sigma Chemical Co. (St. Louis, MO, USA), and all solvents were of HPLC or analytical grade (Merck).

Standard preparation

A stock solution of sesamol (2 µg/mL) was prepared in methanol, which was stable for weeks in the dark and at 0°C. The stock was used for the preparation of working standards (0.05, 0.1, 0.2, 0.3, 0.4, 0.5 µg/mL) and calibration curve.

Sample preparation

Sesame seeds (0.1 g), after grinding in a commercial blender, were extracted with 2 mL of methanol in a shaking incubator at a room temperature. The resulting slurries were centrifuged for 5 min (1000 rpm). The supernatant was kept and the residue was re-extracted under the same condition for three more times. The combined extracts (0.1 g/2500 mL) was injected to HPLC column (100 µL) after filtering through a 0.22 µm filter (Satorius, Goetting, Germany).

Liquid chromatography

The HPLC system consisted of a manual injector (Rheodyne, California, USA), a HPLC pump (Maxi-Star k-1000, Knauer, Berlin, Germany), a C₈ reversed-phase-column [Nova Pak C₈ (4 µm, 250×4.6 mm), Waters, Milford, USA], a spectrophotometer UV-Visible detector (k 2500, Knauer, Berlin, Germany) controlled by a computer software (Eurochrom 2000 ver. 106, Knauer, Berlin, Germany). The mobile phase consisted of methanol. The flow rate was 1 mL/min and the analysis was carried out at room temperature. The UV detector was set at 294 nm and 0.001 AUFS.




Figure 1. HPLC chromatograms of sesamol in: a) Calibration standard solution, b) *karaj 29* cultivar sample

Statistical analysis

All experiments were carried out in triplicate. Mean ± SD was reported for each case and the significance of differences among means were determined at $p < 0.05$ using one way ANOVA followed by Tukey's multiple range test.

Results

We noticed that determination at 294 nm was much more suitable. The chromatograms of sesamol are shown in Figure 1.

Linearity and analytical range

The correlation between the peak area ratios and the sesamol concentrations was evaluated over the range 0-0.5 µg/mL and was found to be linear ($y = 76.063x + 0.5316$; $r^2 = 0.999$; $n = 6$).

Accuracy

The accuracy of the method was verified by

Table 1. precision, sensitivity of method.

Precision (CV%)			
($\mu\text{g/ml}$)	Repeatability (n=3)	Internal reproducibility (n=3)	
Standards:			
0.1 ($\mu\text{g/ml}$)	1.4	2.3	
0.2 ($\mu\text{g/ml}$)	3.5	3.2	
0.4 ($\mu\text{g/ml}$)	2.8	5.9	
Sample (darab 14)	3.0	16.4	
Sensitivity and linearity range			
Calibration (n=6)	Range R ²	Slope (RAUC/ $\mu\text{g/ml}$)	Intercep (RAUC)
0-0.5 $\mu\text{g/ml}$	0.999	76.06	0.5316

means of recovery assay.

This was accomplished by analyzing standard solution and spiked (enriched) sample. The analytical recovery was 100.33% for total sesamol.

Precision

The repeatability of the method was calculated by using the measured data from 3 successive days. Both values were expressed by coefficient variation (CV %). Some method validation (13), (14) data, including precision and linearity, are shown in Table 1. The data of validation method (Table 1) showed that the HPLC is a suitable method for sesamol analysis.

Concentration of sesamol in samples

Sesamol content (mg/g) of Iranian sesame seed brands are shown in Table 2.

Discussion

The mean content of sesamolignan in 7 brands of Iranian sesame seeds (n=63) was 4.67 ± 0.92 mg/g, ranging between 2.75 and 6.13 mg/g (Table 2). Among the 7 brands of Iranian sesame seeds, the samples of *Karaj 29* (5.84 ± 0.25 mg/g), *Dezful* (5.48 ± 0.08 mg/g) and *Varamin* (5.4 ± 0.1 mg/g) had higher concentrations of sesamol, while the sample of *Darab 14*. (3.30 ± 0.57 mg/g) had the lowest ($p < 0.05$).

A few studies in other countries investigated the contents of lignans in sesame seeds or sesame oil. The sesamol content of 14 brands of roasted commercial sesame oils in Taiwan is

reported as 0.30 ± 0.11 mg/g by the total lignans of 11.51 ± 2.81 mg/g, and in Japanese sesame oil, sesamol content was 0.03 ± 0.03 mg/g by the total lignans of 8.02 ± 2.93 mg/g (15). As reported, Indian sesame oil content of the total lignans is 10-20 mg/g. Sesamol was the lowest among the sesame lignans (16).

Sesamol is also produced from the degradation of sesamolignin during the roasting process of sesame seeds (17). Namiki reported that the mean sesamolignin content (1.7 mg/g) was lower than that of Japanese sesame oil (3 mg/g) and the level of sesamol was 10 times higher than that reported in Japan (15). The results were expected because Japanese sesame seeds were not roasted prior to expelling the oil, so sesamol could not be detected in the unroasted seed oil (17). Sesamol presents in crude sesame oil in small amounts (11).

In a study, crude sesame seeds were examined, and determination of sesamol was performed on extract of sesame, rather than on sesame oil. There are two basic strains of sesame seeds: black and white. It was observed that samples were apparently darker, had higher contents of sesamol, so they postulated that the colour of the oils reflected the content of sesamol (18). Nagashima also reported that the water extract (black materials) of black sesame seed coats possessed strong antioxidant activity (19). The total antioxidant status as determined by Trolox equivalent antioxidant capacity assay and expressed as Trolox equivalents was highest for black sesame (65.9 ± 1.7 mg/g), while white seeds showed the lowest (4.4 ± 0.6),

Table 2. Sesamol content of Iranian sesame seeds.

Cultivar	N	Mean (mg/g)	SD	Minimum (mg/g)	Maximum (mg/g)
<i>Branching Naz</i>	9	4.4	0.79	3.51	5.24
<i>Darab 14</i>	9	3.3	0.57	2.75	4.08
<i>Ultan</i>	9	4.07	0.48	3.4	4.65
<i>Non branching Naz</i>	9	4.2	0.54	3.76	5.05
<i>Karaj 29</i>	9	5.84	0.25	5.3	6.06
<i>Varamin</i>	9	5.4	0.1	4.66	6.13
<i>Dezful</i>	9	5.48	0.08	4.42	6.01
Total	63	4.67	0.92	2.75	6.13

concentrated mainly in the hull fraction (20). Otherwise, Mohamed believes that the brown variety contained higher amounts of total sterols and tocopherols but lower amounts of sesamin, sesamol, and total hydrocarbons than the white variety (21). A conflicting report is that higher values of sesamin and sesamol were found in the oil sample from the white sesame seed variety (22). But, in our study, there was no difference between the amount of sesamol content among the seeds of different colours, as the highest content was for *Karaj 29* (white), *Dezful* (dark brown), and *Varamin* (brown). This may indicate that the colour is related to the total antioxidant content, not just sesamol.

Sesamol is a potent phenolic antioxidant (15). Fukuda reported (11) that the addition of 0.5 g/kg sesamol was found to enhance the antioxidative action of γ -tocopherol at all concentrations. Kajimoto found (23) that sesamol had a great preventive effect on the thermal decomposition of tocopherol in oil. But, it seems that this level is insufficient to explain the high stability of the oil, suggesting the presence of other antioxidants. It is important that no single compound can be considered responsible for this antioxidant activity. Combinations of a number of minor constituents such as tocopherols, sesamol, squalene, and antipolymerization sterols in the sesame seed have a synergistic role, increasing the oxidation stability (21, 24-28).

From ancient times to today, sesame has been considered to be a valuable oil-seed, not only because of its medicinal uses, but also because of its medical effects. Some valuable components in sesame contribute to a nutritional and functional

food for humans. Our results indicated that Iranian sesame seeds possess antioxidant components such as sesamol, so the seeds can be easily incorporated into a normal diet at a level that might benefit health as a natural antioxidant with wide food-related applications.

Acknowledgement

The authors acknowledge Tehran University of Medical Sciences for their support through a grant and we thank the Pharmaceutical Sciences Research Center for their help.

References

- (1) Shyu YS and Hwang LS. Antioxidative activity of the crude extract of lignan glycosides from unroasted Burma black sesame meal. *Food Res. Int.* (2002) 35: 357-365
- (2) Salunkhe DK, Charan JK, Adsule RN and Kadam SS. Sesame. In: Van Nostrand R. (ed.) *World Oil Seeds: History, Technology and Utilization*. Spring, New York (1991) 371-402
- (3) Fukuda Y and Namiki M. Recent studies on sesame seed and oil. *Nippon Shokuhin Kogyo Gakkaishi* (1988) 35: 552-562
- (4) Coulman KD, Liu Z, Hum WQ, Michaelides J and Thompson LU. Whole sesame seed is as rich a source of mammalian lignan precursors as whole flaxseed. *Nutrition and Cancer* (2005) 52: 156-165
- (5) Hsu DZ, Su SB, Chien SP, Chiang PJ, Li YH and Lo YJ. Effect of sesame oil on oxidative-stress associated renal injury in endotoxemic rats: involvement of nitric oxide and proinflammatory cytokines. *Shoc.* (2005) 24: 276-280
- (6) Hirata F, Fujita k, Ishikura Y, Hosoda K, Ishikawa T and Nakamura H. Hypocholesterolemic effect of sesame lignan in humans. *Atherosclerosis* (1996) 122: 135-136
- (7) Shimizu S, Akimoto K, Shinmen Y, Kawashima

- H, Sugano M and Yamada H. Sesamin is a potent and specific inhibitor of delta 5 desaturase in polyunsaturated fatty acid biosynthesis. *Lipids* (1991) 26: 512-516
- (8) Lemcke-Norojarvi M, Kamal-Eldin A, Appelqvist LA, Dimberg LH, Ohrvall M and Vessby B. Corn and sesame oils increase serum gamma tocopherol concentrations in healthy Swedish women. *J. Nutr.* (2001) 131: 1195-1201
- (9) Yoshida H and Kajimoto G. Microwave heating affect composition and oxidative stability of sesame (*sesamum indicum*) oil. *J. Food Sci.* (1994) 59: 613-616
- (10) Sonntag N.o.v. Composition and characteristics of individual fats and oils. In: Swern D. (ed.) *Bailey's Industrial Oil and Fat Products*. Wiley, New York (1981) 289-477
- (11) Fukuda Y, Nagata M, Osawa T and Namiki M. Chemical aspects of the antioxidative activity of roasted sesame seed oil and the effect of using the oil for frying. *Agric. Biol. Chem.* (1986) 50: 857-862
- (12) Namiki M. Antioxidants, antimutagens in foods. *Crit. Rev. Food. Sci. Nutr.* (1990) 29: 273-300
- (13) Wood R. *Quality in the Food Analysis Laboratory*. The Royal Society of Chemistry, London (1998) 71-85
- (14) Schwedt G. *The Essential Guide to Analytical Chemistry*. Wiley, New York (1999) 2-16
- (15) Namiki M. The chemistry and physiological functions of sesame. *Food Rev. Int.* (1995) 11: 281-329
- (16) Hemalatha S, Raghunath M and Ghafooruniss A. Dietary sesame oil inhibits iron-induced oxidative stress in rats. *Br. J. Nutr.* (2004) 92: 581-587
- (17) Yoshida H and Takagi S. Effects of seed roasting temperature and time on the quality characteristics of sesame oil. *J. Sci. Food Agric.* (1997) 75: 19- 26
- (18) Wu WH. The contents of lignans in commercial sesame oil of Taiwan and their changes during heating. *Food Chem.* (2007) 104: 341- 344
- (19) Nagashima M, Fukuda Y and Ito R. Antioxidative lignans from industrial waste water in cleaning of black sesame seed. *Nippon Shokuhin Kagaku Kogaku Kaishi* (1999) 46: 1382-1388
- (20) Shahidi F, Liyana-pathirana CM and Wall D. Antioxidant activity of white and black sesame seeds and their hull fractions. *Food Chem.* (2006) 99: 478-483
- (21) Mohamed HMA and Awatif II. The use of sesame oil unsaponifiable matter as a natural antioxidant. *Food Chem.* (1998) 62: 269-276
- (22) Yoshida H, Shigezaki J, Takagi S and Kajimoto G. Variations in the composition of various acyl lipids, tocopherols and lignans in sesame seed oils roasted in a microwave oven. *J. Sci. Food Agric.* (1995) 68: 407-415
- (23) Kajimoto G, Kanomi Y, Kawakami H and Hamatani M. Effects of antioxidants on the thermal oxidation of oils and decomposition of tocopherol in vegetable oils. *J. Jpn. Soc. Nutr. Food Sci.* (1992) 45: 291-295
- (24) Hajimahmoodi M, Oveisi MR, Sadeghi N, Jannat B, Bahaeddin Z and Mansoori S. Gamma tocopherol content of Iranian sesame seeds. *Iranian J. Pharm. Res.* (2008) 7: 135-139
- (25) Suja KP, Abraham JT, Thamizh SN, Jayalekshmy A and Arumughan C. Antioxidant efficacy of sesame cake extract in vegetable oil protection. *Food Chem.* (2004) 84: 393-400
- (26) Chang LW, Yen WJ, Chin Huang S and Duh PD. Antioxidant activity of sesame coat. *Food Chem.* (2002) 78: 347-354
- (27) Lee J, Lee Y and Choe E. Effect of sesamol, sesamin, and sesamolin extracted from roasted sesame on the thermal oxidation of methyl linoleate. *LWT* (2008) in press.
- (28) Suja KP, Jayalekshmi A and Arumughan C. Antioxidant activity of sesame cake extract. *Food Chem.* (2005) 91: 213-219

This article is available online at <http://www.ijpr-online.com>